
The Role of Common Weeds and Insects in the Movement and Severity of Almond Leaf Scorch

Project No.: 07-PATH7-Daane

Project Leader: Kent Daane
Department of Environmental Science
Policy and Management (ESPM)
University of California
Berkeley, CA 94720
(559) 227-9670
daane@ucl.ac.uk

Project Cooperators: Christina Wistrom, ESPM, University of California - Berkeley
Elaine B. Shapland, ESPM, University of California - Berkeley
Glenn Y. Yokota, ESPM, University of California - Berkeley
Roger A. Duncan, University of California Cooperative Extension – Stanislaus County
Joseph H. Connell, University of California Cooperative Extension – Butte County
Mario A. Viveros, University of California Cooperative Extension – Kern County

Interpretive Summary:

Xylella fastidiosa is a xylem-limited bacterium that causes almond leaf scorch (ALS), Pierce's disease (PD) of grapevines, and other plant diseases. To better understand the epidemiology of *X. fastidiosa* and biology of its xylem-feeding insect vectors, vegetation and Cicadomorph insects were surveyed for abundance and *X. fastidiosa* presence in ALS-infected almond orchards in the northern, central, and southern San Joaquin Valley over two years.

Vegetation samples were tested for *X. fastidiosa* by immunocapture PCR. Eleven of 38 species of common ground vegetation tested were positive for *X. fastidiosa*, including shepherd's purse (*Capsella bursa-pastoris*), filaree (*Erodium* spp.), cheeseweed (*Malva parvifolia*), burclover (*Medicago polymorpha*), annual bluegrass (*Poa annua*), London rocket (*Sisymbrium irio*), and chickweed (*Stellaria media*). Sixty-three of 1369 samples tested positive for *X. fastidiosa*, with *Xylella* detection between November and March.

Four point nine percent (4.9%), of more than 42,000 Cicadomorph insects collected were xylem feeders, overwhelmingly green sharpshooters, *D. Minerva*. Insect populations were highly seasonal, with few vectors collected from December to mid-

April, increasing in late April, and peaking from mid-June through mid-July. Green sharpshooters were eliminated inside the orchard when vegetation was removed for almond harvest in mid to late July, although they were still found in adjacent intact vegetation into September. Sites in Butte and Glenn counties had much higher green sharpshooter populations (2.49 and 8.77 *D. minerva*/100 sweeps), than sites in Stanislaus (0.66 *D. minerva*/100 sweeps), or Kern counties (0.08 *D. minerva*/100 sweeps). *D. minerva* were collected more frequently at the margins (8.4/100 sweeps), compared to >10m inside (1.5/100 sweeps), and > 10m outside the almond orchards (3.3/100 sweeps). There were no differences in green sharpshooter populations on riparian habitat, on orchard floor vegetation, and on weeds near roadsides. One point one percent (1.1%), of green sharpshooters tested were positive for *X. fastidiosa*, which were collected between May and July inside the almond orchard. *X. fastidiosa* strains in sharpshooters matched strains isolated from ALS-infected trees and weeds.

Both ground vegetation and almond trees were most commonly infected with the almond strain of *X. fastidiosa*. ALS-infected almond samples had a *X. fastidiosa* concentration within previously reported ranges (1.84×10^6 - 2.15×10^7 CFU/g), and between 1.9 and 0.17% of almond trees at the study sites had ALS in 2004.

An insecticide spray applied to almond orchard floor vegetation in January at one of the study sites did not suppress green sharpshooter populations or populations of Cicadomorphs in general for the following growing season.

Our results show that the biology of the green sharpshooter, the main ALS vector, is much different from the biology of the blue-green sharpshooter or the glassy-winged sharpshooter, the main PD vectors. Green sharpshooters feed and breed on common ground vegetation species that can harbor *X. fastidiosa* on the almond floor. While the proportion of almond trees, insects, and alternate weedy hosts with *X. fastidiosa* is low, green sharpshooter populations can be high, and ground cover extensive in orchards with flood irrigation or with abundant winter rainfall. In orchards with ALS-susceptible varieties, sampling to determine ALS prevalence and green sharpshooter occurrence, and a year-round weed control program may assist in management of *X. fastidiosa*.

The Study Objectives Were:

- To characterize the *X. fastidiosa* presence and strains found in almond leaf scorch samples and compare them with strains from nearby alternate host plants, especially almond orchard floor vegetation.
- To collect insects in ALS-infected orchards and nearby vegetation, and the conduct laboratory analyses determine whether or not they carry *X. fastidiosa*.
- To test *D. minerva* for *X. fastidiosa* transmission efficiency to and from almonds and weeds commonly found in almond orchards.
- Collect regional data on ALS epidemiology with respect to orchard management (e.g., irrigation) and the surrounding environmental conditions (e.g., nearby crop plantings).
- To describe native sharpshooter biology and development.

- To conduct experiments to reduce native sharpshooter density during winter and spring periods.

Materials and Methods:

Sample sites were selected based on grower reports of ALS incidence, and located in the northern (ca. 122°09'W, 39°39'N and 121°50'W, 39°45'N, Glenn County; ca. 121°56'W, 39°49'N and 121°55'W, 39°36'N Butte County), central (ca. 121°03'W, 37°31'N and ca. 120°44'W, 37°39'N, Stanislaus County), and southern Central Valley (ca. 119°17'W, 35°22'N and ca. 119°15'W, 35°20'N, Kern County). Vegetation and insects were collected every 2 to 6 weeks, depending on seasonal abundance.

Vegetation Sampling:

On each visit, researchers walked through the orchard and conducted a visual survey for the four most abundant weed species. Vegetation was identified according to the Statewide IPM Program produced by the Agriculture and Natural Resources Department at the University of California, Davis (available online). Four evenly spaced rows were then selected (300 – 400 m per row), and 3 to 5 leaves of each of the common weed species were collected from each transect. Samples were stored separately for each species and transect in a 3.8-liter plastic bag. Ten to thirty individual plants were sampled for each weed species in each transect. The collected material was stored in a cooler (ca. 7°C), and processed for *X. fastidiosa* presence within 2 days of collection.

Insect Sampling:

Each orchard was divided into transects based on the surrounding environment (i.e. the presence of a creek, alfalfa field, or other potential breeding site of *D. minerva*). Orchard floor vegetation in each transect was swept 100 times with a 35 cm diameter sweep net, repeated three times in each transect, approximately 1200 sweeps per site per sampling date. Five to ten yellow sticky traps (7.5cm by 13cm), were set in lower branches of almond trees, 1 to 2m above the ground, in the center and around the perimeter of each site. Insects collected by sweep net and sticky trap were stored at -20 °C until sorting. Xylem-feeding Cicadellid and Cercopid Homopterans were identified and counted at each site and sampling date, along with Delphacid and Deltacephalinid leafhoppers, and Membracids. A subset of collected *D. minerva*, *X. fulgida*, and *S. festinus* were preserved in acetone (Fukatsu 1999), for later detection of *X. fastidiosa* in mouthparts. Specimens were identified according to Bland (1978), and Dietrich (2005), and representative vouchers submitted to the Essig Museum of Entomology at U.C. Berkeley.

Bacterial Detection in Vegetation:

Each sample (plant species and transect), was processed separately for the presence of *X. fastidiosa* using immunocapture DNA separation, which is a sensitive antibody-linked system that selectively removes bacteria (Shapland 2006). We selected plant

parts where sharpshooters were known to feed, such as leaf petioles, leaf blade bases of grasses, and plant stems. Briefly, 0.70g of plant material was cut into 1-mm sections, placed into 3 ml of modified SCP buffer, and thoroughly homogenized (Brinkmann Instruments, Westbury NY). The homogenate was strained through cheesecloth into a sterile microcentrifuge tube and centrifuged at 14000x g for 5 min. The pellet was resuspended in 500 µl PBS/BSA buffer and 100 µl of rabbit antibody to *X. fastidiosa* (diluted 1/100 in PBS/BSA), was added to each sample. Samples were incubated at room temperature for 30 min with gentle shaking, followed by centrifugation at 14000x g for 2.5 min. The pellet then was resuspended in 1.0 ml of PBS/BSA, centrifuged a second time for 2.5 min, and resuspended in 1.0 ml of PBS/BSA. Five µl of Dynabeads M-280 sheep anti-rabbit IgG (DynaL Biotech, Lake Success NY), was added to each sample and incubated at room temperature for 30 min with gentle shaking. The beads were removed with a Magnetic Particle Concentrator (DynaL MPC-S, Oslo), and washed three times with 1.0 ml, 1.0 ml, and 500 µl of PBS/BSA. Finally, the bead-antibody-bacteria complex was resuspended in 10 µl of sterile distilled water.

For PCR amplification, all samples were prepared in sterile microcentrifuge tubes with 12.5 µl Taq Master Mix (Qiagen, Valencia, CA), 6.5 µl PCR water, 1 µl of each primer RST-31 and RST-33, and 4 µl of DNA extract. PCR reactions were carried out in a thermal cycler according to the conditions described by Minsavage, et al. (1994). PCR products were separated by electrophoresis in 1.5% agarose gel, stained with ethidium bromide and viewed under ultraviolet light. The presence of *X. fastidiosa* in the original sample was determined by a band at 733 kb.

After PCR amplification, strain differences were detected with *RsaI* restriction enzyme digestion (Hendson et al. 2001). *RsaI* cuts PCR products from oak, oleander, peach, plum, and all but three ALS strains into two fragments, while leaving PD strains intact.

Bacterial Detection in Insects:

X. fastidiosa was detected in insects utilizing vacuum-extraction and PCR (Bextine 2005). Insect specimens were removed from the acetone; the heads were removed, and pinned with a #2 insect pin, avoiding the eyes. The pinned head was placed into a 96-well plate filled with 180µL of cell lysis buffer (ATL) from the DNeasy tissue kit for animal tissues (Qiagen). The plate was placed into a vacuum chamber which was pressurized and depressurized five times to flush the lytic buffer through the insect head. The ATL buffer was then processed according to the manufacturer's directions. The extracted DNA was amplified by PCR with RST 31-33 primers as described above. The *X. fastidiosa* strain present in insects was identified with multiplex PCR (Hernandez-Martinez 2006) or restriction enzyme digestion (Hendson 2001), and compared with *X. fastidiosa* strains prevalent in ALS-infected trees in the orchard (Shapland 2006).

Mapping and Detection of Almond Leaf Scorch:

All field sites were surveyed for the presence and distribution of ALS. Disease symptoms were rated visually at Butte, Glenn and Stanislaus county sites in September and October 2004 and October 2005, and in November 2004 in Kern County. The locations of diseased trees were recorded with a GPSMAP 76 handheld device (Garmin Inc., Olathee KS). All trees at Grainland Rd., Zeering Rd., Tim Bell Rd. and Kern sites 2, 6 and 7 were surveyed, while subsets of trees were surveyed at Rd. P, Rd. VV, and Anita Rd.

Bacterial Titer and Strain:

X. fastidiosa was cultured from symptomatic almond trees at each site and from fresh samples of alternate host plants at sites where previously collected samples tested positive for *X. fastidiosa*, using immunocapture DNA separation and PCR amplification. In late summer and early fall, when the symptoms of ALS were most obvious we surveyed each orchard and collected symptomatic almond leaf and petiole samples. Both ground vegetation and almond samples were processed for bacterial culture on selective media within 24h. We plated samples on both PWG and PD3 media to provide a rudimentary indication of *X. fastidiosa* strain. Grape strains of *X. fastidiosa* grow on both PWG (Davis et al. 1983), and PD3 (Davis et al. 1981), but some almond strains grow only on PWG (Almeida and Purcell 2003). Samples were cultured (Hill and Purcell 1995), to determine both the strain and concentration of bacteria in almond and ground vegetation samples.

D. Minerva Biology and Development:

D. minerva colonies were established with insects collected from Stanislaus (near Modesto), Sonoma (near Guerneville), and Solano (near Fairfield), counties, and maintained in the greenhouse at UC Berkeley. Insects were raised on 'sweet' basil (*Oncimum basilicum*), and bermudagrass (*Cynodon dactylon*,) plants in 3.75 L pots enclosed by wood and mesh cages. The greenhouse was maintained at 25°C (+10 °C /-5°C), with natural lighting, supplemented from October to April with plant and aquarium lights (General Electric, Fairfield, CT), for a 14-hour day length. All plants were grown in supersoil (Rod McLellan Co., Marysville, OH), and fertilized daily with diluted 20-20-20 fertilizer.

Seeds of burclover (*Medicago polymorpha*), cheeseweed (*Malva parvifolia*), johnsongrass (*Sorghum halepense*), and filaree (*Erodium* sp.), were collected in the field and germinated in the greenhouse. Basil was grown from purchased seeds, and bermudagrass was vegetatively propagated by stolons. Plants were used in experiments when they were six to nine weeks old. Scientific and common names of the weeds were listed according to the Jepson Manual (Hickman 1993), and the Growers Weed Identification Handbook (Anonymous 1999), and voucher specimens deposited at the Jepson Herbarium at U.C. Berkeley. In all experiments, equal numbers of each plant species were set up in each replication to control for the effects of

greenhouse temperature, plant nutritional status, light exposure on the maturation and behavior of *D. minerva*, and the effects of plant age, phenology and nutrition.

Nymph Development:

Ten *D. minerva* adults, five males and five females, were transferred to bermudagrass, burclover, cheeseweed, and filaree for a 48-hour egg laying period, contained in a cylindrical plastic cage with mesh screen. Adults were removed after two days; the plant was re-caged, and monitored for nymph emergence thereafter. Approximately 10 insects were gently shaken off each plant weekly, and the numbers of each instar were counted. Insects were returned to the plant after counting, and adult sharpshooters were removed from the plant as they matured. All insects were moved to a new plant of the same species as plants became senescent or infested with other insects. Experiments were conducted from June to November 2006, with eleven replications of equal numbers of plants from each species per replication.

Adult Survival:

To determine the suitability of the various weed species for long-term *D. minerva* survival, eight newly-moulted adults (<14 days old), four males and four females, were placed on 'Peerless' almond seedlings, bermudagrass, burclover, cheeseweed, seedling 'Cabernet Sauvignon' grapes, johnsongrass, filaree and basil. Eight replications with equal numbers of plants from each species per replication were completed from February to July 2007. Every week, all adults were gently shaken off the plant and counted, then returned to the plant. All insects were moved to a new plant of the same species when a plant was senescent or infested with other insects.

Egglaying:

Five adult male, and five adult female *D. minerva* were transferred to bermudagrass, burclover, cheeseweed, johnsongrass, filaree and basil in seven replications with equal numbers of plants from each species per replication, from February through May 2007. Adults were removed after a four-day egglaying period, and the eggs present in the plant visualized plant with a Nikon SMZ800 dissecting microscope. Eggs and plants were photographed with a Nikon Coolpix digital camera. The plant was returned to the insectary, and re-caged. The numbers of nymphs emerging from the plant were counted 7, 14 and 21 days after egglaying. Nymphs were removed from the plants after counting. Once all nymphs were removed, plant dry weights were determined by removal of plant material at the soil line, which was placed in a paper bag, dried for at least seven days in a heated cabinet, and weighed.

X. Fastidiosa Survival in, and D. Minerva Transmission Between Almonds and Alternate Hosts:

Tested between almonds and nine weed species (Shapland et al 2006): shepherd's purse (*Capsella bursa-pastoris*), filaree, cheeseweed, burclover, common groundsel (*Senecio vulgaris*), London rocket (*Symbrium irio*), sowthistle (*Sonchus* sp.), chickweed (*Stellaria media*), and burning nettle (*Urtica urens*). Plants were grown in the greenhouse from seed as previously described.

Mechanical Inoculation of Alternate Hosts:

X. fastidiosa isolates (ALS6 and Fresno-ALS), were started 10 to 14 days prior to inoculations. *X. fastidiosa* colonies from one plate were suspended in 2 ml SCP buffer by pipetting repeatedly to break up clumps of bacteria. Inoculum concentrations were determined by dilution plating on PWG media. Plants were inoculated immediately after preparing the bacterial suspension, always on a sunny day to ensure the drop was drawn into the plant via transpiration. A 5 µl drop of *X. fastidiosa* suspension was placed on the inoculation location and probed 3 to 6 times with a #2 insect pin. Sites were marked with tape.

Plants were assessed for *X. fastidiosa* presence and titer four weeks after inoculation. Two samples were cut immediately adjacent to the inoculation site: one for culturing and the other was frozen at -20C for immunocapture PCR (Shapland 2006), if needed due to contamination from bacteria other than *X. fastidiosa*.

Also, twenty plants of each species were inoculated as above and planted outside in screenhouses in the field, half in Bakersfield CA, and half in Parlier, CA. Plants were assessed for *X. fastidiosa* presence four, six and eight weeks after mechanical inoculation via culture and immunocapture PCR as previously described.

Insect Inoculation of Alternate Hosts:

Almond plants infected with *X. fastidiosa* were generated by mechanically inoculating almonds with ALS6 or Fresno-ALS strains two to three months prior to transmission test, as previously described. Almonds and weeds were kept in unsprayed greenhouse until symptoms developed, then tested for *X. fastidiosa* presence and titer via culture two weeks prior to use in transmission experiments.

Between 80 and 100 green sharpshooters were placed on *X. fastidiosa*-infected almonds for a 4-day acquisition access period (AAP). *D. minerva* were then moved in groups for four to alternate hosts for a four-day inoculation access period (IAP), confined to plants in small foam or mesh cages. Following inoculation, inoculation site was marked with lab tape and plants were placed in the greenhouse to develop infections.

Insect Acquisition of *X. Fastidiosa* from Alternate Hosts:

Alternate hosts belonging to the nine weed species were infected with *X. fastidiosa* by mechanical inoculation with ALS6 and Fresno-ALS as previously described. Plants were four to six weeks old when inoculated. *D. minerva* free of *X. fastidiosa* were reared through three weekly changes as nymphs on basil.

D. minerva were placed in groups of ten on basil plants for a four-day pretest period, then divided into two for transfer to alternate hosts for a four-day AAP, then transferred in those groups to seedling 'Peerless' almonds for a four-day IAP. At the end of the IAP, insect heads were preserved in acetone and vacuum-extraction PCR performed to detect the presence of *X. fastidiosa* in *D. minerva*. Ten to 12 weeks after inoculation, the almond seedlings were tested for *X. fastidiosa* infection by culture.

Dormant Spray to Suppress Vector Populations:

The treated orchard was located southwest of Chico (ca.121°55'W, 39°36'N, Butte County), 20.2 hectares of mixed Peerless, Nonpareil, and Price varieties, between 19 and 26 years old. In surveys in fall 2004 and 2005, ALS-infected trees were present, predominantly in the center of the orchard, but also in individual trees in the eastern and northern edges. Except for the insecticide spray, standard cultural practices were followed. Vegetation was removed from the orchard in preparation for nut harvest on 15 July 2005, ending insect sampling on the orchard floor vegetation.

The orchard was divided into 8 blocks of 460 trees per block. The groundcover on the almond orchard floor in four of the blocks was treated with Asana XL (esfenvalerate, DuPont Corp. Wilmington, DL), insecticide at 10 oz. per acre (0.725 L per hectare), on 27 January 2005. Sites were sampled for Cicadomorph insects, particularly *D. minerva*, once a month from February through July 2005 as previously described. One hundred sweeps per block on the orchard floor vegetation were taken at each sampling date.

Data Analysis:

Data were entered in Excel (Microsoft Corp., Redmond WA), with averages compared using one-way and repeated measures ANOVA, and means separated using Tukey's HSD comparison (JMP, SAS Institute, Cary NC and Systat, San Jose CA).

Results and Discussion:

Vegetation Sampling:

From June 2003 to April 2005, 38 species of ground vegetation were commonly found (Table 1, as presented in the 2005 report), with most material collected in winter and spring when ground vegetation was abundant. Between August and October, it was difficult to find live ground vegetation within the almond orchards, as the orchard floor was completely free of vegetation prior to harvest. Irrigation was also discontinued, so

that the nuts can be shaken from the trees and dried on the bare ground. All orchards in this study followed these practices. At that time samples were across the orchard rows (transects), to produce a single sample for each plant species and orchard (Figure 1, as presented in the 2005 report).

Insect Sampling:

From June 2003 to September 2005, more than 42,000 Cicadomorph insects were collected in Stanislaus, Butte, Kern and Glenn counties, including 1,920 *D. minerva*, five *X. fulgida*, one *P. spumarius*, and 80 *S. festinus*. Four point nine percent (4.9%), were xylem feeders (sharpshooters or spittlebugs), known vectors of *X. fastidiosa* (Figure 2). *X. fulgida* was only collected at the Anita Rd. site, and only in sweeps of riparian vegetation along a creek adjacent to the almond orchard. Other Cicadellid species collected in Stanislaus and Butte counties included *Exitanus exitosus* Uhler, and *Ambleysellus grex* Oman. *Thamnotettix zelleri* Kirshbaum was collected from grasses at Anita Rd. in May. *Actinopterus* sp., *Empoasca* sp., and *Exitanus exitosus* Uhler were collected in Kern County. *Nothodelphax consimilis* Van Duzee (Delphacidae), also was collected at all field sites in Butte, Glenn and Stanislaus counties, but not at the Kern County sites.

Insect populations were highly seasonal, with few vectors collected from December to mid-April, increasing populations in late April, and peak populations from mid-June through mid-July. Cicadellid populations were eliminated inside the orchard when vegetation was killed with herbicides in preparation for almond harvest in mid to late July, although *D. minerva* were still found in adjacent riparian areas or crops into September.

More *D. minerva*, and Cicadomorphs in general were found at northerly collection sites (Figure 2). The highest Cicadomorph populations were at Anita Rd. in Butte county (146 per 100 sweeps), and lowest were populations at Zeering Rd. in Stanislaus county (7 per 100 sweeps). Sites in Butte and Glenn counties had higher *D. minerva* populations (17.0, 2.9, 2.3, to 1.3 *D. minerva* per 100 sweeps), than the Stanislaus county sites (1.1 and 0.4 *D. minerva* per 100 sweeps). In Kern County, *D. minerva* were only found at Site 6 in June 2004, 1.3 per 100 sweeps). The exception to this trend was *S. festinus* populations, as 55 *S. festinus* were collected in Kern County, 25 were collected at Butte and Glenn county sites, and none were collected from sites in Stanislaus County.

Four thousand eighty-four insects, including nine *D. minerva*, were collected by sticky traps hung in the lower branches of the trees for 13,051 trap-days, compared to 42,392 collected by sweeps (including 1,912 *D. minerva*). No insects were collected in limited sampling with beat sheets at four sites (0 in 1660 beats), compared to 1,318 insects collected in 430 sweeps at the same sites and dates.

More sharpshooters were collected in sweeps at the margins of orchards (an average of 8.4 per 100 sweeps), compared to collections more than 10m inside (1.5 per 100 sweeps) and more than 10m outside the almond orchards (3.3 per 100 sweeps) at the

Chico and Modesto-area study sites (Sharpshooters: one-way ANOVA, $P = 0.0001$, $n = 983$). There were no significant differences between total Cicadomorph populations at the three collection locations (Total Cicadomorphs: one-way ANOVA, $P = 0.17$, $n = 983$; Figure 3).

There were no differences in sharpshooter populations between collections made on riparian habitat, almond orchard floor vegetation, and roadside vegetation, for collections made on the same days and at the same study sites (one-way ANOVA, $P > 0.27$, $n = 217$). Total numbers of Cicadomorphs were lower in riparian habitats compared to roadside or almond orchard floor vegetation. (One-way ANOVA, $P = 0.07$, $n = 217$; Figure 4).

Bacterial Detection in Vegetation:

Sixty-three of 1,369 samples from the six orchards were positive for *X. fastidiosa* (4.6%). *X. fastidiosa* was recovered from 11 of the 38 ground vegetation plant species tested, including *Capsella bursa-pastoris* (shepherd's purse), *Malva parvifolia* (cheeseweed), *Senecio vulgaris* (common groundsel), *Sisymbrium irio* (London rocket), *Sonchus* sp. (sowthistle), *Stellaria media* (chickweed), *Urtica urens* (burning nettle), and *Veronica persica* (speedwell), eight species from which it had not previously been recovered in the field (Table 1). There was a strong seasonal component to bacterial presence in ground vegetation, with no *X. fastidiosa* positive samples found between April and mid-October during the two years of the study (Table 2, as presented in the 2005 report). Linear regression showed a positive, significant relationship between the number of samples taken per plant species and the percentage of samples positive for *X. fastidiosa* ($y = 0.0553x - 0.2074$, $R^2 = 0.8935$).

Restriction enzyme digestion of PCR products and bacterial culture on selective media showed that almond trees at six of seven studied orchards were infected with the almond strain of *X. fastidiosa* (Table 3, as presented in the 2005 report). At Zeering Rd. in Stanislaus County, a grape strain of *X. fastidiosa* was isolated from all weeds and almond trees sampled. At each site, tissue samples from both almond trees and surrounding weeds gave the same result: each contained either the grape or almond strain of *X. fastidiosa*, but never both.

Bacterial Detection in Insects:

X. fastidiosa was recovered from 17 of 1,532 (1.1%), of sharpshooters tested. Three *D. minerva* tested positive at Anita Rd., 13 at Rd. VV, and one at Tim Bell Rd. Only almond strain *X. fastidiosa* was recovered from sharpshooters, matching strains isolated from ALS-infected trees and weeds at the Anita Rd., Rd. VV, and Tim Bell Rd. sites. *X. fastidiosa*-positive sharpshooters were collected between May 19 and July 24 in the Chico area and on April 4 in Stanislaus County. All sharpshooters testing positive for *X. fastidiosa* were collected inside the almond orchard.

Mapping and Detection of X. Fastidiosa Infection in Almond Trees:

Overall infection percentages were low, between 1.9 and 0.17% at the study sites in 2004 (Table 3). More ALS-symptomatic trees were observed in 2004 than 2005, when trees had lost many leaves due to senescence and *Tranzchelia discolor* almond rust. All sites included highly susceptible varieties of trees: Sonora, Solano, or Peerless. At Rd. VV in Glenn County, no *X. fastidiosa*-infected trees were detected; however, *X. fastidiosa* was detected in weeds and insects at that site.

Bacterial Titer and Strain:

Petioles from ALS-symptomatic almond trees at Zeering Rd. containing the grape strain of *X. fastidiosa* had an average concentration of 2.15×10^6 CFU/g, which was significantly greater than the concentrations at other sites sampled ($P = 0.014$). Our results agreed with previous findings that the average *X. fastidiosa* titer in ALS-symptomatic almond leaves was lower than the average *X. fastidiosa* titer in PD symptomatic grapes (Almeida and Purcell 2003).

Nymph Development:

D. minerva nymphs developed from egg to adulthood on all species tested: burclover, bermudagrass, cheeseweed, and filaree (Figure 5). The longest maturation time recorded was 70 days on bermudagrass, and 49 days for the other three species. The shortest maturation time was 28 days, on all plants. Although a few nymphs emerged 7 days after egg laying, most emerged after 14 days. *D. minerva* appeared to mature at similar rates on all weed species tested. Seventy-nine percent (348 of 441), of sharpshooters survived the 2-day egg laying period, similar proportions on all plants (Chi-square with pairwise comparisons, $P > 0.10$).

Egg laying:

Numbers of emerging nymphs were similar on all five plant species tested (one-way ANOVA, $P = 0.52$), with lots of variability in numbers of nymphs between replications (Figure 6). Overall, the most nymphs emerged from johnsongrass, (19 average per plant), followed by bermudagrass.

D. minerva laid eggs in masses in lower stems and petioles of weed species tested, often quite close to the soil line. Eggs were deposited under the plant epidermis, completely covered by plant tissue (Figure 7, A and B). Tests are planned for fall 2007 to determine the number of eggs per mass, and egg laying habit on the various plant species.

D. minerva survived the four-day egg laying period (8.88 adults per plant, SE = 0.20), in similar proportions on all five plant species (one-way ANOVA, $P = 0.93$). There were no differences in dry weight between species of plants used for egg laying, with average dry weights of 3.0, 2.9, 2.6, 2.1 and 1.9g per plant, for cheeseweed, filaree, johnsongrass, burclover, and bermudagrass, respectively (one-way ANOVA, $P = 0.37$). There was no

correlation between plant dry weight and numbers of nymphs from that plant (simple linear regression, $R^2 = 0.009$, $P = 0.57$).

The average temperature in the insectary during the egg-laying interval for replications six through ten was between 22.9 and 25.0°C. High temperatures measured from 38.7 to 36.9°C, and low temperatures from 18.4 to 20.6°C. There was no apparent relationship between total nymph emergence and average, maximum high or maximum low temperatures.

Adult Survival:

D. minerva adults survived longest on Johnsongrass, averaging 48 days, and shortest on almonds, an average of 26 days. Other species were as follows: basil, 37 days; bermudagrass, 35 days; burclover, 39 days; cheeseweed 40 days; and grapes, 33 days. For all plant species tested, at least one insect survived the full 70 day test period (Figure 8).

X. fastidiosa Survival in, and *D. Minerva* Transmission Between Almonds and Alternate Hosts:

D. minerva transmitted *X. fastidiosa* from almonds to four of the nine weed species tested in the greenhouse: cheeseweed, groundsel, London rocket and sowthistle. *X. fastidiosa* populations were similar in groundsel (5.1×10^6 CFU/g), London Rocket (7.5×10^5 CFU/g), and sowthistle (6.2×10^5 CFU/g). Bacterial populations were significantly lower in cheeseweed (4.4×10^3 CFU/g; one-way ANOVA with log log₁₀-transformed data; Tukey-Kramer HSD test, $P = 0.003$).

Both grape and almond strains of *X. fastidiosa* reached similar populations in weeds. ALS6 populations averaged 2.8×10^5 CFU (colony-forming-units) per gram of plant material, and Fresno-ALS averaged 8.9×10^5 CFU/g in infected weeds (t-test with log₁₀-transformed data; $P = 0.15$).

Mechanical Inoculation of Alternate Hosts:

X. fastidiosa infection rates and titers were lower in inoculated alternate hosts grown in the field compared to the same plants grown in the greenhouse (Table 4). The experiments were completed in August 2008, and data analysis is ongoing. However, preliminary analysis suggests that *X. fastidiosa* populations were roughly 10 times lower in field-grown plants. The plant species from which *X. fastidiosa* was detected regularly in surveys of almond orchard vegetation (Table 2), appear to be frequently-infected, high-titer hosts when mechanically inoculated, except for shepherd's purse. Experiments to test insect inoculation of *X. fastidiosa* into, and acquisition from, these nine alternate host species were finished in summer 2007.

Dormant Spray to Suppress Vector Populations:

There were no statistically significant differences in total Cicadomorph or *D. minerva* populations between insecticide-treated and untreated blocks (Figure 9). Overall, mean

Cicadomorph populations were greater in treated blocks (77.6 per 100 sweeps), compared to untreated blocks (58.4 per 100 sweeps), while *D. minerva* populations were lower in sprayed blocks (1.2 per 100 sweeps), than in unsprayed blocks (0.7 per 100 sweeps).

Previous field surveys for *X. fastidiosa* in alternate host plants focused on Pierce's disease management in surveys of common weed and riparian vegetation species inside and adjacent to vineyards. With the recent increase of ALS in California, there was need to identify *X. fastidiosa* hosts in almond orchard floor vegetation as well. Different vegetation types are present near grapes and almonds, as well as different vector species. In grapes, the blue-green sharpshooter (*Graphocephala atropunctata*), or glassy-winged sharpshooters (*Homalodisca vitripennis*), are the primary vectors. In almonds, the green sharpshooter, *Draeculacephala atropunctata*, is the main vector. The three species have very different life cycles and patterns of host plant use. There are also different vegetation patterns in vineyards and almond orchards. In vineyards, a clear edge effect was found with PD incidence closely tied to adjacent riparian habitat (Purcell and Frazier 1985), whereas most previous work did not show any clear spatial patterns with ALS in almond orchards (Purcell 1980).

Sharpshooters on grapes, especially the blue-green, generally enter vineyards from adjacent vegetation, feed briefly on grapevines, and do not breed or persist inside vineyards. In contrast, almond orchards, especially in areas with abundant winter rainfall or flood irrigation, often contain *D. minerva* breeding and feeding habitat through mid-summer. Adult longevity, egg laying, and nymph development studies conducted in the greenhouse indicated that *D. minerva* feed, reproduce and mature on weed species that are *X. fastidiosa* hosts and commonly found inside almond orchards. Thus the vectors of ALS multiply inside the orchard, in contrast to PD vectors which mostly originate outside the vineyard. The distribution of *D. minerva* may explain differences ALS and PD distribution. We are currently conducting transmission tests in the greenhouse, to determine the rates of *X. fastidiosa* transmission by *D. minerva* between the most frequently-infected alternate hosts and almonds.

D. minerva were eliminated inside the orchard when vegetation was killed with herbicides in preparation for almond harvest in mid to late July, although they were still found in adjacent riparian areas or crops into September. Sites in northerly counties had much higher *D. minerva* populations than sites in central or southern San Joaquin valley, indicating that *D. minerva* are highly dependant on vegetation present in the orchard, as opposed to the almond trees themselves, or migrating into the orchard from outside.

Four point six percent (4.6%), of field-collected, naturally-infected vegetation samples contained *X. fastidiosa*. The *X. fastidiosa*-positive plant species were the most-frequently found species in the orchards on most of the sampling dates. There was a positive and significant relationship between the number of samples taken per plant species and the percentage of samples positive for *X. fastidiosa*. We detected *X. fastidiosa* in weeds between October and April, whereas insect populations were highly

seasonal, with few vectors collected from December to mid-April, increasing in late April, and peaking from mid-June through mid-July.

One point one percent (1.1%), of *D. minerva* tested were positive for *X. fastidiosa*. *X. fastidiosa*-positive sharpshooters were collected between May and July. The seasonal difference between when *X. fastidiosa* was detected in weeds (November through April), and when *X. fastidiosa* was detected in insects may be explained two ways:

1. Since the overall proportion of *D. minerva* with *X. fastidiosa* was so low, infected sharpshooters were only detectable in the months when populations were highest.
Or
2. The orchard floor vegetation was in better condition and temperatures more favorable to *X. fastidiosa* growth and survival in plants, in the winter and early spring. Since adult *D. minerva* survived at least 10 weeks on weed species present in the orchard in greenhouse longevity tests, it is possible that a sharpshooter could acquire *X. fastidiosa* in early spring and be detected when collected in the summer months.

Both ground vegetation and almond trees were most commonly infected with the almond strain of *X. fastidiosa*. *X. fastidiosa* strains in sharpshooters matched strains isolated from ALS-infected trees and weeds. While a low proportion of almond trees at the study sites had ALS, those trees were highly symptomatic, and had *X. fastidiosa* populations between 10^6 and 10^7 colony-forming-units per gram of petiole – thus, infected trees were infrequent but had lots of inoculum sufficient for sharpshooter acquisition.

While data analysis is still ongoing as of August 2008, *D. minerva* transmitted *X. fastidiosa* from almonds to four weed species in greenhouse tests: cheeseweed, groundsel, London rocket and sowthistle. Of the four, sowthistle was the most frequently infected (35% of inoculated plants became infected), and is the most persistent in the field. There were no differences between the ability of “grape” (Fresno-ALS), and “almond” (ALS6), strains of *X. fastidiosa* to infect alternate hosts, and both bacterial strains reached similar populations in weedy alternate hosts. Overall, 55% of weeds inoculated using needle-inoculation developed *X. fastidiosa* infections, compared to only 23% of plants inoculated with *D. minerva*. Studies that rely upon mechanical inoculation exclusively thus may overestimate the potential for weeds or other hosts to support reservoir populations of *X. fastidiosa*.

Pyrethroid insecticide treatment, applied to orchard floor vegetation in late January, was not effective in reducing ALS vector populations or total populations of leafhoppers and Delphacids. One reason for the lack of effectiveness may be the timing of the application, as the insecticide was applied four months before peak *D. minerva* populations. The very low populations at the first sampling date in February (and none in March and April), show that the few adults that survived the winter multiply to yield a much larger second generation in May and June. This population pattern was found in other almond orchards in Butte, Glenn, and Stanislaus counties, and in previously-published studies in pastures and alfalfa fields in Fresno County. With an estimated half-life of 7.5 days in lab tests (Anonymous 2000), Asana insecticide may also not

persist long enough to effectively target emerging *D. minerva* in late and mid-February. The application of insecticide in late January may have been too early to kill overwintering *D. minerva* adults as they came out in February. Finally, *D. minerva* may have been physically protected from the spray, since they overwinter hidden in perennial vegetation. Future experiments should determine the most effective timing, the best cultural or chemical techniques to control *D. minerva* populations, and to determine an economic threshold for *D. minerva* infestation.

Our results show that the biology of ALS vectors is different from PD vectors, and that they feed and breed on common ground vegetation species that can harbor *X. fastidiosa* on the almond floor. While the proportion of almond trees, insects, and alternate weedy hosts with *X. fastidiosa* is low, *D. minerva* populations can be high, and ground cover extensive in orchards with flood irrigation or in areas with abundant winter rainfall. In orchards with ALS-susceptible varieties, sampling to determine ALS and *D. minerva* occurrence, and a year-round weed control program may assist in management of *X. fastidiosa*.

References:

- Almeida, R. P. P. and Purcell, A. H. 2003. Biological traits of *Xylella fastidiosa* strains from grapes and almonds. *Appl. Environ. Microbiol.* 69:7447-7452.
- Anonymous. 1999. The Grower's Weed Identification Handbook. Univ. Calif. Div. Agric. Natural Resources Publication No. 4030.
- Anonymous. 2002. Asana Insecticide Technical Bulletin: <http://www2.dupont.com>. DuPont Corp., Wilmington, DL.
- Bextine, B., Blua, M., Harshman, D., and TA Miller. 2005. A SYBR green-based real-time PCR protocol and novel DNA extraction technique to detect *Xylella fastidiosa* in *Homalodisca coagulata*. *J. Econ. Entomology* 98: 667-672.
- Bland, R.G. 1978. How to know the Insects, 3rd Edition. WCB/McGraw-Hill Publishers, New York, NY. 156-166.
- Davis, M. J., Whitcomb, R. F., and Gillaspie, A. G. J. 1981. Fastidious bacteria of plant vascular tissue and invertebrates (including so-called rickettsia-like bacteria). 1230-1234. in: *The Prokaryotes: A Handbook on Habits, Isolation, and Identification of Bacteria*. Springer-Verlag, Berlin, Heidelberg.
- Davis, M.J. Raju, B.C., Brlansky, R.H., Lee, R.F. Timmer, L.W., Norris, R.C., and McCoy, R.E. 1983. Periwinkle wilt bacterium: axenic culture, pathogenicity, and relationships to other gram-negative, xylem-inhabiting bacteria. *Phytopathology* 73: 1510-1515.
- Dietrich, C.H. 2005. Keys to the families of Cicadomorpha and subfamilies and tribes of Cicadellidae (Hemiptera: Auchenorrhyncha). *Florida Entomol.* 88: 502-517.
- Fukatsu, T. 1999. Acetone preservation: a practical technique for molecular analysis. *Molecular Ecology* 8: 1935-1945.
- Hendson, M., Purcell, A. H., Chen, D., Smart, C., Guilhabert, M., and Kirkpatrick, B. 2001. Genetic diversity of Pierce's disease strains and other pathotypes of *Xylella fastidiosa*. *Appl. Environ. Microbiol.* 67:895-903.

- Hernandez-Martinez, R., Costa, H.S., Dumenyo, C.K., and D.A. Cooksey. 2006. Differentiation of strains of *Xylella fastidiosa* infecting grape, almonds, and oleander using a multiprimer PCR assay. *Plant Disease* 90: 1382-1388.
- Hickman, J.C (ed). 1993. *The Jepson Manual: Higher Plants of California*, 2nd ed. University of California Press, Berkeley, CA.
- Hill, B. L. and Purcell, A. H. 1995. Multiplication and movement of *Xylella fastidiosa* within grapevine and four other plants. *Phytopathology* 85:1368-1372.
- Minsavage, G. V., Thompson, C. M., Hopkins, D., Leite, R. M. B. V. C., and Stall, R. E. 1994. Development of a polymerase chain reaction protocol for detection of *Xylella fastidiosa* in plant tissue. *Phytopathology* 84:456-461.
- Purcell, A.H. 1980. Almond leaf scorch: leafhopper and spittlebug vectors. *J. Econ. Entomol.* 73: 834-838.
- Purcell, A.H. and N.W. Frazier. 1985. Habitats and dispersal of the principal leafhopper vectors of Pierce's disease bacterium in the San Joaquin valley. *Hilgardia* 53: 1-32.
- Shapland, E.B., Daane, K.M., Yokota, G.Y., Wistrom, C., Connell, J.H., Duncan, R.A., and M.A. Vivieros. 2006. Ground vegetation survey for *Xylella fastidiosa* in California almond orchards. *Plant Disease* 90: 905-909.

Recent Publications:

Shapland, E.B., Daane, K.M., Yokota, G.Y., Wistrom, C., Connell, J.H., Duncan, R.A., and Vivieros, M.A. 2006. Ground vegetation survey for *Xylella fastidiosa* in California almond orchards. *Plant Disease* 90: 905-909.

A second manuscript, *Seasonal Abundance and Natural Infectivity of Xylella fastidiosa Vectors in California Almond Orchards*, is in preparation for submission to the *Journal of Economic Entomology*.

Additional presentations of research results were at the CDFA GWSS-PD Research Symposium in 2005 and 2006, and at the 2006 Almond Board Conference in Modesto, CA and the 2007 ESA Conference in Portland, OR.

Table 1: Ground Vegetation Collected from Almond Orchards and Tested for *Xylella Fastidiosa*.

Weed species		<i>X. fastidiosa</i>		Reference
Scientific Name	Common Name	This study	Previous studies	
<i>Amaranthus spp.</i>	Pigweed	-	-	6
			+	36*
<i>Amsinckia spp.</i>	Fiddleneck	-	-	6
<i>Anagallis arvensis</i>	Scarlet pimpernel	-	-	6
<i>Avena fatua</i>	Wild oat	-	+	11
<i>Brassicaceae spp.</i>	Mustards	-	+	6
<i>Capsella bursa-pastoris</i>	Shepherd's purse	+	N/A	
<i>Chamaesyce maculata</i>	Spotted spurge	-	N/A	
<i>Chamomilla suaveolens</i>	Pineapple weed	-	N/A	
<i>Chenopodium album</i>	Lambsquarter	-	N/A	
<i>Claytonia perfoliata</i>	Miner's lettuce	-	+	30,31
<i>Conyza bonariensis</i>	Fleabane	-	N/A	
<i>Conyza canadiensis</i>	Horseweed	-	+	36*
<i>Coronopus didymus</i>	Lesser swinecress	-	N/A	
<i>Cyperus esculentus</i>	Yellow nutsedge	-	+	11
<i>Echinochloa crus-galli</i>	Barnyard grass	-	+	17*, 36*
<i>Erodium spp.</i>	Filaree	+	+	6, 11*, 36*
<i>Escallonia montevidensis</i>	Escallonia	-	+	11
<i>Festuca spp.</i>	Fescue grass	-	N/A	
<i>Helianthus spp.</i>	Sunflower	-	+	6*
<i>Hordeum murinum</i>	Hare barley	-	+	11
<i>Lactuca serriola</i>	Prickly lettuce	-	+/-	30,31, 36*
<i>Malva parvifolia</i>	Cheeseweed	+	-	6, 11*
<i>Medicago polymorpha</i>	Burclover	+	+	11
<i>Poa annua</i>	Annual bluegrass	+	+	11
<i>Portulaca oleracea</i>	Common purslane	-	-	11*, 36*
<i>Ranunculus spp.</i>	Buttercup	-	N/A	
<i>Rumex crispus</i>	Curly dock	-	+	11
<i>Salsola tragus</i>	Russian thistle	-	N/A	
<i>Senecio vulgaris</i>	Common groundsel	+	N/A	
<i>Sisymbrium irio</i>	London rocket	+	N/A	
<i>Sonchus oleraceus</i>	Annual sowthistle	-	+	36*
<i>Sonchus spp.</i>	Sowthistle	+	-	6
<i>Sorghum halepense</i>	Johnsongrass	-	+/-	11*, 30, 31
<i>Stellaria media</i>	Chickweed	+	N/A	
<i>Typha spp</i>	Cat tail	-	N/A	
<i>Urtica urens</i>	Burning nettle	+	N/A	
<i>Veronica persica</i>	Speedwell	+	N/A	
<i>Xanthium strumarium</i>	Cocklebur	-	+	36*

Results are compared to previous field surveys near vineyards and riparian areas, except for those marked * which refer to greenhouse studies.

Table 2: *X. Fastidiosa* Detection in Almond Orchard Vegetation. Data Combined from Six Almond Orchards, from June 2003 to April 2005.

Plant species	J	F	M	A	M	J	J	A	S	O	N	D
Burning nettle		■										
Common Groundsel											■	
Speedwell		●										
Sowthistle			■									
London Rocket		■	■									
Annual bluegrass			■									●
Cheeseweed	◆		■ ●							◆	■	◆ ●
Chickweed		●	●		■						●	●
Burclover			◆									
Shepherd's purse		●	◆		■						●	
Filaree			◆ ♠ ●							◆	■ ●	●

■ = Kern, ● = Stanislaus, ◆ = Butte, ♠ = Glenn

Table 3: Almond Orchards Surveyed for Cicadomorph Insects, Vegetation, and *Xylella Fastidiosa* Strains and Titters.

Site (County)	GPS Coordinates	No. trees surveyed	No. trees XF +	Strain in almonds	Strain in weeds	<i>X. fastidiosa</i> titer ²
Grainland Rd. (Butte)	121°55'W, 39°36'N	3700	7	Almond	Almond	1.84 x 10 ⁶ ab
Anita Rd. (Butte)	121°56'W, 39°49'N	~6000	38	Almond	Almond	2.30 x 10 ⁶ b
Rd. P (Glenn)	122°09'W, 39°39'N	~3600	6	Almond	Almond	1.19 x 10 ⁷ ab
Rd. VV (Glenn)	121°50'W, 39°45'N	~2500	0	--	Almond	--
Zeering Rd. (Stanislaus)	121°03'W, 37°31'N	2560	49	Grape	Grape	2.15 x 10 ⁷ a
Tim Bell Rd. (Stanislaus)	120°44'W, 37°39'N	900	2	Almond	Almond	6.27 x 10 ⁶ b
Site 2 (Kern)	-	-	4	Grape	--	3.90 x 10 ⁶
Site 6 (Kern)	-	-	3	Almond	--	1.84 x 10 ⁷
Site 7 (Kern)	-	-	14	Almond	Almond	1.08 x 10 ⁷ b

¹ Results based on PCR and restriction enzyme digestion with *Rsa1*.

² Ave. CFU/g cultured from ALS symptomatic petioles: 27 July, 30 Sept. and 14 October 2004.

³ Different letters after each mean indicate a significance difference (P < .05), Tukey's pairwise comparison. Kern sites 2 and 6 not included due to small number of sampled trees.

Table 4: *X. Fastidiosa* Infection in Greenhouse (GH) and Field-Grown Alternate Hosts Following Needle Inoculation with ALS6 and Fresno-ALS Strains.

Infections were assessed after four to eight weeks in field-grown plants, and four weeks in greenhouse-grown plants. Results are preliminary for greenhouse-grown plants, with anticipated completion in fall 2007.

Species	Growing location	No. tested	No. infected (%)	CFU/g
Burclover - <i>Medicago polymorpha</i>	GH	36	21 (58)	1.77 x 10 ⁷
	field	54	13 (24)	3.00 x 10 ⁶
Cheeseweed - <i>Malva parviflora</i>	GH	-	-	-
	field	38	6 (16)	1.35 x 10 ⁷
Chickweed - <i>Stellaria media</i>	GH	35	30 (86)	5.93 x 10 ⁷
	field	5	3 (60)	-
Filaree - <i>Erodium</i> sp.	GH	-	-	-
	field	6	2 (33)	6.44 x 10 ⁷
Common groundsel - <i>Senecio vulgaris</i>	GH	-	-	-
	field	28	3 (11)	3.34 x 10 ³
London rocket - <i>Sysimbrium irio</i>	GH	17	8 (47)	1.61 x 10 ⁶
	field	17	0 (0)	-
Burning nettle - <i>Urtica urens</i>	GH	27	9 (33)	1.09 x 10 ⁶
	field	14	1 (7)	-
Shepherd's purse - <i>Capsella bursa-pastoris</i>	GH	-	-	-
	field	2	0 (0)	-
Sowthistle - <i>Sonchus oleraceus</i>	GH	50	15 (30)	3.87 x 10 ⁷
	field	38	3 (8)	8.33 x 10 ⁴

Figure 1: *Xylella Fastidiosa* Presence in Vegetation in Almond Orchards. Combined Results from 6 Almond Orchards in Butte, Glenn, Stanislaus, and Kern Counties from June 2003 to April 2005.




Figure 2: Average Cicadomorph and Sharpshooter Populations from June 2003 to September 2005 inside and adjacent to A) Four Almond Orchards near Chico, CA (Butte and Glenn Counties), and B) Two Almond Orchards near Modesto, CA (Stanislaus County).




Figure 3: Average Insect Populations Collected from Sweeps of Orchard Floor and Adjacent Vegetation >10m Outside, Within 10m of the Margins, and >10m on the Interior of Study Sites.




Figure 4: Average Insect Populations Collected from Sweeps of Vegetation on the Almond Orchard Floor (Almond), in Adjacent Riparian Habitat (Riparian), and on Roadsides Adjacent to Study Sites. Insects were Collected on the Same Days at the Same Study Sites.




Figure 5: *D. Minerva* Maturation on Four Species, in Average Number of Insects per Plant, on Alternate Hosts in Seventy Ddays in the Greenhouse: A) Filaree, B) Cheeseweed, C) Burclover, and D) Bermudagrass.




Figure 6: *D. Minerva* Nymph Emergence from Five Weed Species in Greenhouse Tests. Eight Adult *D. Minerva* were Given a 4-Day Egg Laying Interval on Alternate Host Plants of Similar Age and Size; Adults were Removed, and Emerged Nymphs Counted Weekly Thereafter.




Figure 7: *D. Minerva* Egg Masses in Johnsongrass Approximately 10 Days after Laying. A. Dissected to Remove Flap of Olanth Epidermis and Show Developing Eggs; B. Intact Egg Mass.




Figure 8: Green Sharpshooter Adult Survival on Eight Host Plant Species in Greenhouse Tests.




Figure 9: Total Cicadomorph (A: top) and *D. Minerva* (B: bottom) Populations Swept from Vegetation in a Butte County Almond Orchard in 2005. Treatment was with Asana Insecticide on 27 January 2005. Means of the Two Treatments are not Significantly Different in Analysis with Repeated Measures ANOVA (Cicadomorphs: $P = 0.25$, $n = 47$, *D. Minerva*: $P = 0.68$, $n = 47$).

