

---

## Field Evaluation of Almond Varieties

---

**Project No.:** 14-HORT2-Lampinen

**Project Leader:** Bruce Lampinen  
Department of Plant Sciences  
University of California, Davis  
One Shields Ave.  
Davis, CA 95616  
530.752.2588  
bdlampinen@ucdavis.edu

**Project Cooperators:** Tom Gradziel, Sam Metcalf, William Stewart, Maria Contador, and Mary Ann Thorpe, UC Davis  
Joe Connell (emeritus), UCCE - Butte County  
Dani Lightle, UCCE – Butte/Glenn/Tehama Counties  
Roger Duncan, UCCE - Stanislaus County  
Gurreet Brar, UCCE - Fresno/Madera Counties  
Mario Viveros and Minerva Gonzalez, UCCE - Kern County  
Chico State University  
The Billings Ranches, Kern County

**Objectives:**

The first objective of this project is to evaluate performance of pollenizers and Nonpareil clones in a replicated field trial in McFarland California. The second objective is to use the replicated data on light interception and yield to assess the relative productivity per unit of light intercepted for the Nonpareil clones, varieties and selections. The third objective is to initiate the next generation almond variety trials.

**Interpretive Summary:**

Yields at the McFarland trial continue to be high compared to the 1993 regional almond variety trials. The McFarland trial reached an average production of 3000 kernel pounds per acre in the 5<sup>th</sup> growing season which was six years earlier than the 1993 Kern RAVT while neither the 1993 Butte nor the 1993 Delta trials ever reached this level of production. In order to develop the canopy this quickly, water and other inputs need to be high and this has resulted in high disease pressure (especially hullrot). Because this trial is replicated, unlike the earlier regional almond variety trials, we have the opportunity to more accurately assess yield differences among the Nonpareil clones, varieties and selections. In addition, because we have detailed light interception data, we can assess whether differences in productivity among varieties and selections are due to faster canopy growth or higher productivity per unit light intercepted. Nonpareil yields varied from 4300 to almost 5000 kernel pounds per acre in 2011 and continued to be generally higher than pollenizer yields. In 2012, yields were down substantially compared to 2011, most likely due to alternate bearing. In 2013, yields increased but were not as high as 2011. Yields in 2014 were still quite high. There appears to be

some separation of cumulative yields for the Nonpareil clones with Nonpareil-Nico highest and Nonpareil-5, Nonpareil-6 and Nonpareil-Jones all having significantly lower cumulative yields. Hull rot continued to be the major disease related problem with the highest levels in 2-19e and Kochi with the Nonpareil clones having intermediate hull rot pressure. Hull rot pressure was minimal in 2013 and 2014. Summary data from the 2006 to 2014 seasons shows that although bloom period varied significantly from one year to the next, in general, overlap of Nonpareil bloom with pollenizers was good in all years.

The next generation of Regional Almond Variety Trials was planted in the winter of 2014 in Butte (Chico State University), Stanislaus (Salida School District Site), and Madera (Chowchilla grower site) counties. Nonpareil was alternated with 30 varieties and/or selections at all 3 sites. Trees at the Butte, Stanislaus and Madera trial were planted on Krymsk 86, Nemaguard and Hansen 536 rootstocks respectively (with the exceptions listed at the bottom of **Table 5**). Unlike the previous generation Regional Almond Variety Trials, there are four replications of each of the varieties and selections at each of the three sites in the 2014 trials.

## **Materials and Methods:**

### McFarland Variety Trial

This report will concentrate on a replicated variety trial of eight varieties and eight Nonpareil clones that was planted in 2004 in Kern County near McFarland. Soils at the site consist of McFarland loam and Wasco sandy loam (both Class I soils). The irrigation system is double line drip. Tree spacing is 20 feet between tree rows and 18 feet between trees for a density of 121 trees per acre. Varieties planted included Chips, Kahl, Kochi, Marcona, Selection 2-19e, Sweetheart and Winters. Nonpareil clones planted include Nonpareil 3-8-2-70, Nonpareil 5, Nonpareil 6, Nonpareil 7, Nonpareil Dr., Nonpareil-J, Nonpareil-Newell and Nonpareil-Nico. There are six replications of each variety and Nonpareil clone with 34 trees per replication. Pollenizer and Nonpareil rows alternate in the orchard.

In 2014, data was collected on bloom and maturity timing, disease pressure (scab, *Alternaria* and hull rot), yield and midday canopy light interception. In 2006, 2007 and 2008, midday canopy light interception was measured manually using a hand lightbar. The mobile platform light bar (described in report for Almond Board project titled “Development and Testing of a Mobile Platform for Measuring Canopy Light Interception and Stress in Almond”) was used to measure midday canopy light interception at the McFarland trial in 2009-2014.

### 2014 Regional Almond Variety Trials

The next generation almond variety trials were planted in the winter of 2014 in Butte (Chico State University), Stanislaus (Salida School District Site), and Madera (Chowchilla grower site) counties. The varieties and selections planted are listed in **Table 5**. The first 31 items are common to all 3 sites and a few different items added at individual sites are listed at the bottom of **Table 5**. Trees at the Butte, Stanislaus and Madera trial were planted on Krymsk 86, Nemaguard and Hansen 536 rootstocks

respectively (with the exceptions listed at the bottom of **Table 5**). Trees were planted at a spacing of 18' x 22' at the Butte site (110 trees/acre), 16' x 21' at the Stanislaus site (130 trees/acre) and 12' x 21' at the Madera site (173 trees/acre). This density is significantly higher than the previous generation RAVTs where planting densities for the Butte, San Joaquin and Kern trials were 64, 75 and 86 trees/acre respectively.

## Results and Discussion:

### McFarland Trial

The grower started ripping every other row center in 2009 and this has resulted in significant improvements in water penetration in the orchard. This procedure has continued and appears to be beneficial with much better water penetration observed.

Weather during bloom was generally good at the McFarland trial site. For the 2014 bloom period there were 146 good bee flying hours (temperature  $\geq 59^{\circ}\text{F}$ , windspeed  $\leq 10$  MPH, and no rain). This compared to 182, 91, 58, 94, 103, and 175 good bee flying hours in 2008, 2009, 2010, 2011, 2012 and 2013 respectively.

Although the absolute date of bloom for all varieties varied from year to year, bloom overlap among the Nonpareil clones and pollenizers was good in every year (**Figure 2, 3**). Although the bloom dates vary considerably from year to year, the relative bloom dates among the Nonpareil versus pollenizers all appear to shift together. Average bloom dates for the 2006 to 2014 seasons are shown in **Figure 4**.

Hullsplit progression or 2010-2013 is shown in **Figure 5**. In 2014 hullsplit initiation and completion was earlier than average by about 2.5 weeks (**Figure 6**). This is the earliest we have observed during this trial to date. The average hullsplit progression for 2006 to 2014 is shown in **Figure 7**.

In 2010, scab was only observed on Winters and Nonpareil-5 and in 2011 on Winters and selection 2-19e (**Table 1**). In 2012, scab was present at low level on most varieties and selections. In 2013, scab was present only in Kahl, Marcona and Winters and in 2014 only in Kochi and Winters (**Table 1**). *Alternaria* was not present in 2010 but in 2011 and 2012 all Nonpareil selections and pollenizers had some present with the worst being in Kahl, Winters, Kochi and Marcona (**Table 1**). In 2013, *Alternaria* was not observed in Nonpareil selections but was present in Sweetheart, Chips, Kahl, Marcona and Winters (**Table 1**). In 2014, *Alternaria* was worst in Winters, Kahl, Sweetheart and Marcona (**Table 1**). Hullrot continued to be the main problem at this site. There was substantial hullrot in all varieties and selections with the most severe disease pressure in Kochi and Winters in both 2010 and 2011 (**Table 1**). Kochi also displayed severe hullrot in 2008 and 2009. There was also substantial hull rot in all the Nonpareil clones in 2010 and slightly less in 2011 (**Table 1**). Overall, hull rot levels decreased from 2012 to 2014, with 2014 showing the lowest presence in five years (**Table 1**).

Yield data for the McFarland trial from the 2006 through 2014 seasons are shown in **Table 2, 3 and 4**. All of the Nonpareil clones have tended to have higher cumulative yield compared to most of the pollenizers (**Table 4**). Among the Nonpareil clones, there is some separation regarding cumulative yield with Nonpareil Nico, Nonpareil 38270

and Nonpareil Newell having the highest cumulative yields (**Table 4**). Selection 2-19e (Kester) has had the highest cumulative yield among all of the pollenizers in the trial followed by Winters (**Table 4**). The replication provided in this trial adds much value to the data compared to earlier unreplicated variety trials but it comes at a cost in terms of the time required for maintaining, rating and harvesting the plots.

The yields from this trial continue to be high for the age of the orchard. **Figure 1** shows average yield by orchard age for all varieties, selections and Nonpareil sources for the Butte, Delta and Kern trials planted in 1993 compared to the McFarland trial planted in 2004. The McFarland trial reached an average production of 3000 kernel pounds per acre six years earlier than the 1993 Kern RAVT while neither the Butte nor Delta trials ever reached this level of production (**Figure 1**). Yields have tended to alternate around 3700 kernel pounds per acre for the past few years which is what would be expected based on light interception levels.

The productivity per unit canopy light interception was quite high for the size of the tree in 2009 (ranging from 52 to 85 kernel pounds per unit light intercepted) and was still fairly high in 2010 (ranging from 23 to 51 kernel pounds per unit light intercepted) since our previous data has suggested 50 kernel pounds per unit light intercepted is near the normal maximum for almond (**Table 3**). In 2011, yield per unit light intercepted averaged 74.8 for the Nonpareil clones and 40 to 65 for the pollenizers (**Table 3**). This trend towards yield per unit PAR intercepted in the range of 50 has continued in 2013 and 2014 (**Table 4**). **Figure 8** shows the yield per unit PAR intercepted by variety or selection for all years of the study (data for Nonpareil sources are averaged). This figure suggests that the yields tend to alternate around 50 confirming our earlier observations. Nonpareil has produced above the 50 line while the clones have averaged below the line.

#### Next Generation Almond Variety trials

The Butte, Stanislaus and Madera sites were planted in the winter of 2014 as described in the Materials and Methods section. See **Table 5** for a list of items planted at the trials. Bloom, hullsplit and yield data collection will begin in 2016. All of the sites are growing well and although the Chico trial was short about 60 trees from the nursery at planting and some trees were lost at all 3 sites, they have mostly been replaced.

#### Acknowledgements:

The authors wish to thank the Almond Board of California for their continued support of this project. We also want to thank the Billings Ranches for their excellent cooperation in managing and maintaining this trial. In addition, we would like to thank Chico State University/Jeff Boles, the Salida School District/Lane Parker as well as the Creekside Farming Company for their cooperation on the next generation variety trials.

**Table 1.** Scab rating, *Alternaria* rating, and hull rot strikes per tree for the 2010-2014 seasons at the McFarland trial. Letters indicate significant difference at the 5% level of significance.

		Scab Rating	Alternaria rating	Hull Rot Strikes
2010	selection 2-19e	0.00 a	Chips	0.00 a
	Chips	0.00 a	Kahl	0.00 a
	Kahl	0.00 a	Kochi	0.00 a
	Kochi	0.00 a	Marcona	0.00 a
	Marcona	0.00 a	Nonpareil 3-8-2-70	0.00 a
	Nonpareil 3-8-2-70	0.00 a	Nonpareil-5	0.00 a
	Nonpareil-6	0.00 a	Nonpareil-6	0.00 a
	Nonpareil-7	0.00 a	Nonpareil-7	0.00 a
	Nonpareil-DR	0.00 a	Nonpareil-DR	0.00 a
	Nonpareil-J	0.00 a	Nonpareil-J	0.00 a
	Nonpareil-Newell	0.00 a	Nonpareil-Newell	0.00 a
	Nonpareil-Nico	0.00 a	Nonpareil-Nico	0.00 a
	Price	0.00 a	Price	0.00 a
	Sweetheart	0.00 a	selection 2-19e	0.00 a
	Nonpareil-5	1.00 a	Sweetheart	0.00 a
	Winters	2.00 b	Winters	0.00 a
				Winters
2011				539.67 c
	Chips	0.00 a	Nonpareil-Nico	0.67 a
	Kahl	0.00 a	Nonpareil-7	0.67 a
	Kochi	0.00 a	Nonpareil-J	0.83 a b
	Marcona	0.00 a	Sweetheart	1.00 a b
	Nonpareil 3-8-2-70	0.00 a	Nonpareil-Newell	1.00 a b c
	Nonpareil-5	0.00 a	Nonpareil-5	1.00 a b c
	Nonpareil-6	0.00 a	Nonpareil-6	1.00 a b c
	Nonpareil-7	0.00 a	Nonpareil 3-8-2-70	1.00 a b c
	Nonpareil-DR	0.00 a	2-19E	1.17 a b c d
	Nonpareil-J	0.00 a	Nonpareil-DR	1.17 a b c d
	Nonpareil-Newell	0.00 a	Chips	1.50 b c d
	Nonpareil-Nico	0.00 a	Marcona	1.67 c d e
	Sweetheart	0.00 a	Kochi	1.83 d e
	2-19E	0.17 b	Winters	2.33 e
	Winters	3.00 c	Kahl	2.33 e
				Kochi
2012				325.83 d
	Nonpareil-5	0.0 a	Nonpareil-6	0.67 a
	Chips	0.0 a	Nonpareil-Newell	0.83 a
	Kahl	0.2 a b	Nonpareil 3-8-2-70	0.83 a
	Nonpareil-J	0.2 a b	Nonpareil-Nico	0.83 a
	Nonpareil-6	0.2 a b	Nonpareil-DR	0.83 a
	2-19E	0.3 a b c	Chips	1.00 a
	Nonpareil-Nico	0.3 a b c	Nonpareil-J	1.00 a
	Nonpareil-7	0.5 a b c	Sweetheart	1.00 a
	Nonpareil-DR	0.5 a b c	Nonpareil-7	1.00 a
	Marcona	0.5 a b c	Kahl	1.17 a
	Kochi	0.5 a b c	Nonpareil-5	1.17 a
	Nonpareil 3-8-2-70	0.7 b c	Marcona	1.50 b
	Nonpareil-Newell	0.8 c	Kochi	1.50 b
	Sweetheart	1.8 d	2-19E	1.50 b
	Winters	3.0 e	Winters	2.50 c
				2-19E
2013				81.00 g
	Nonpareil-DR	0.0 a	Nonpareil-J	0 a
	Nonpareil-7	0.0 a	Nonpareil-DR	0 a
	Nonpareil-J	0.0 a	Nonpareil-7	0 a
	Nonpareil-5	0.0 a	Nonpareil-Newell	0 a
	Nonpareil-6	0.0 a	Nonpareil-Nico	0 a
	Nonpareil-Newell	0.0 a	Nonpareil 3-8-2-70	0 a
	Kochi	0.0 a	Nonpareil-5	0 a
	Nonpareil 3-8-2-70	0.0 a	Nonpareil-6	0 a
	2-19E	0.0 a	Kochi	0 a
	Sweetheart	0.0 a	2-19E	0 a
	Nonpareil-Nico	0.0 a	Sweetheart	1 a
	Chips	0.0 a	Chips	1 b
	Kahl	0.2 a b	Kahl	1 b
	Marcona	0.3 b	Marcona	1 b
	Winters	1.2 c	Winters	2 c
				2-19E

	<b>Scab Rating</b>		<b>Alternaria rating</b>		<b>Hull Rot Strikes</b>	
2014	Kester (2-19E)	0 a	Nonpareil-J	0.0 a	Kahl	0 a
	Nonpareil-7	0 a	Nonpareil-5	0.0 a	Marcona	0 a
	Nonpareil-Newell	0 a	Nonpareil 3-8-2-70	0.0 a	Sweetheart	1 a
	Nonpareil-J	0 a	Nonpareil-7	0.0 a	2-19E	2 a
	Nonpareil-5	0 a	Nonpareil-Nico	0.0 a	Nonpareil 3-8-2-70	2 a
	Nonpareil 3-8-2-70	0 a	Nonpareil-6	0.0 a	Winters	2 a
	Nonpareil-DR	0 a	Nonpareil-Newell	0.2 a b	Chips	3 a
	Nonpareil-6	0 a	Kochi	0.3 a b	Nonpareil-Nico	3 a b
	Marcona	0 a	2-19E	0.5 b c	Nonpareil-7	4 a b c
	Kahl	0 a	Nonpareil-DR	0.5 b c	Nonpareil-6	5 a b c
	Sweetheart	0 a	Chips	0.5 b c	Nonpareil-Newell	5 a b c
	Nonpareil-Nico	0 a	Marcona	0.8 c d	Nonpareil-5	6 a b c
	Chips	0 a	Sweetheart	1.0 d e	Nonpareil-J	11 b c
	Kochi	0.17 a	Kahl	1.3 e	Nonpareil-DR	11 c
	Winters	2.17 b	Winters	3.0 f	Kochi	25 d

**Table 2.** Number of nuts per tree, average kernel weight, shelling percentage, and kernel pounds per unit of photosynthetically active radiation (PAR) intercepted, kernel pound per acre, cumulative kernel pounds per acre for the 2006 through 2008 seasons. Data for each year is sorted by cumulative yield. Letters indicate significant difference at the 5% level of significance.

Variety	No. of nuts/tree	Average kernel wt	Shelling percentage	Kernel pounds per			Cumulative kernel yield (lbs/acre)
				unit PAR int.	Tree	Acre	
2-19e	6852 a	0.94 g	53.0 d	45.5 a	14.2 a	1718 a	1718 a
Winters	6648 a	0.87 h	53.4 d	47.9 a	12.7 a	1540 a	1540 a
Marcona	3611 bcd	1.31 a	30.7 f	41.4 a	10.4 b	1258 b	1258 b
Nonpareil-Ni	4246 b	1.09 cde	67.2 a	32.4 b	10.2 bc	1232 bc	1232 bc
Nonpareil-5	3713 bcd	1.12 bcd	67.9 a	30.1 b	9.2 bcd	1110 bcd	1110 bcd
Nonpareil-D	3867 bc	1.07 def	63.4 abc	29.9 b	9.1 bcd	1103 bcd	1103 bcd
Nonpareil-3-8-2-70	3848 bc	1.07 cde	64.6 ab	30.6 b	9.1 bcd	1101 bcd	1101 bcd
Nonpareil-Ne	3815 bc	1.07 cde	67.7 a	28.2 b	9.0 bcd	1086 bcd	1086 bcd
Nonpareil-6	3886 bcd	1.12 bc	67.0 a	28.3 b	8.9 bcd	1075 bcd	1075 bcd
Nonpareil-J	3717 bcd	1.08 cde	64.0 abc	28.4 b	8.8 bcd	1066 bcd	1066 bcd
Chips	3623 bcd	1.02 f	53.8 d	26.9 b	8.1 bcde	985 bcde	985 bcde
Kochi	3134 cd	1.16 b	59.9 c	26.7 b	8.0 cde	965 cde	965 cde
Nonpareil-7	3288 bcd	1.08 cde	65.1 a	24.4 bc	7.8 de	941 de	941 de
Kahl	3139 cd	1.06 ef	47.8 e	31.0 b	7.4 de	889 def	889 def
Sweetheart	2777 d	0.95 g	67.8 a	16.7 c	5.6 f	681 f	681 f
Variety	No. of nuts/tree	Average kernel wt	Shelling percentage	Kernel pounds per			Cumulative kernel yield (lbs/acre)
				unit PAR int.	Tree	Acre	
2-19e	13149 a	0.78 e	54.3 d	58.4 ab	22.8 a	2756 a	4474 a
Winters	11972 ab	0.83 de	60.2 b	65.8 a	21.8 ab	2634 ab	4173 a
Nonpareil-Newell	10659 bc	0.90 bc	67.3 a	52.9 bc	21.0 abc	2543 abc	3626 b
Nonpareil-Nico	9260 cde	0.92 bc	66.0 a	47.1 cde	18.8 abcd	2279 abcd	3511 b
Nonpareil-Driver	9793 cd	0.91 bc	65.6 a	51.1 bc	19.6 abcd	2371 abcd	3474 b
Nonpareil-5	8905 cdef	0.95 b	67.0 a	49.3 bcd	19.0 abcd	2299 abcd	3410 b
Nonpareil-3-8-2-70	9340 cde	0.92 bc	66.3 a	50.8 bc	18.9 abcd	2291 abcd	3393 b
Nonpareil-7	9517 cd	0.92 bc	67.9 a	48.1 bcde	19.3 abcd	2332 abcd	3272 bc
Marcona	6938 fg	1.08 a	29.8 f	52.8 bc	16.5 def	1995 def	3252 bcd
Kahl	9594 cd	0.91 bc	47.6 e	65.2 a	19.3 abcd	2332 abcd	3222 bcd
Nonpareil-J	9137 cde	0.89 bcd	65.5 a	44.4 cde	17.8 bcde	2152 bcde	3218 bcd
Nonpareil-6	8396 def	0.94 b	67.1 a	43.3 cde	17.4 cde	2103 cde	3178 bcd
Chips	7681 defg	0.87 cd	54.4 d	38.9 def	14.7 ef	1780 ef	2766 bcd
Kochi	6006 g	1.08 a	59.4 bc	37.9 ef	14.3 ef	1729 ef	2694 de
Sweetheart	6767 fg	0.89 bcd	66.6 a	31.2	13.1 f	1588 f	2165 e
Variety	No. of nuts/tree	Average kernel wt	Shelling percentage	Kernel pounds per			Cumulative kernel yield (lbs/acre)
				unit PAR int.	Tree	Acre	
2-19e	13472 a	0.93 g	54.3 d	56.5 bcde	27.4 cd	3321 cd	7795 a
Nonpareil-Nico	13879 a	1.10 cd	66.0 a	67.5 a	33.5 a	4056 a	7567 ab
Nonpareil-3-8-2-70	12506 bcd	1.17 cd	66.3 a	66.0 a	30.7 b	3714 b	7106 bc
Nonpareil-5	12883 ab	1.08 de	67.0 a	63.9 ab	30.5 b	3692 b	7102 bc
Nonpareil-Newell	11916 bcd	1.09 de	67.3 a	57.3 bcde	28.6 cd	3456 cd	7086 bc
Nonpareil-Driver	12729 abc	1.07 de	65.6 a	62.5 abc	29.8 bc	3611 bc	7085 bc
Nonpareil-7	13250 ab	1.06 de	67.9 a	62.3 abc	31.1 ab	3763 ab	7035 bc
Winters	9872 e	1.02	60.2 b	53.4 def	22.1 fg	2670 g	6843 c
Nonpareil-6	10707 de	1.16 c	67.1 a	54.7 cde	27.3 cd	3300 cd	6478 cd
Nonpareil-J	11071 d	1.09 cde	65.5 a	54.6 cde	26.6 de	3224 de	6442 cd
Kahl	10720 de	0.96 fg	47.6 e	61.2 abcd	22.6 fg	2733 fg	5954 de
Chips	11465 cd	0.97 fg	54.4 d	51.8 ef	24.4 ef	2956 ef	5722 ef
Sweetheart	13149 ab	0.82 g	66.6 a	45.3 f	23.9 ef	2893 ef	5059 g
Marcona	4721 f	1.39 a	29.8 f	36.9	14.5 h	1748 h	5001 g
Kochi	5882 f	1.28 b	59.5 bc	35.0	16.5 h	2002 h	4696 g

**Table 3.** Number of nuts per tree, average kernel weight, shelling percentage, and kernel pounds per unit of photosynthetically active radiation (PAR) intercepted, kernel pound per acre, cumulative kernel pounds per acre for the 2009 through 2012 seasons. Data for each year is sorted by cumulative yield. Selection 2-19e was recently released as Kester.

2009		Variety	No. of nuts/tree	Average kernel wt (g)	Shelling percentage	Kernel pounds per			Cumulative kernel yield (lbs/acre)
unit PAR int.	Tree	Acre							
Nonpareil-Nico	13773 ab		1.05 bcd	74.7 ab	69.3 abcd	31.8 ab	3851 a	11417 a	
Nonpareil-Newell	14513 a		1.03 bcd	74.8 ab	72.8 abc	33.1 a	4004 a	11090 a	
2-19e	14706 a		0.84 f	65.6 f	71.6 abc	27.1 c	3285 c	11080 a	
Nonpareil-Driver	13856 ab		1.08 ab	75.8 a	76.1 ab	32.9 a	3977 a	11062 a	
Nonpareil-3-8-2-70	13756 ab		1.04 bcd	74.6 ab	71.8 abc	31.4 ab	3798 ab	10905 abc	
Nonpareil-7	13051 ab		1.03 bcd	72.6 abc		29.5 bc	3571 bc	10606 abc	
Nonpareil-5	12070 bcd		1.08 ab	74.2 ab	68.5 abcd	28.7 bc	3476 bc	10577 abc	
Nonpareil-6	13505 ab		1.02 bcd	71.2 cd	68.9 abcd	30.2 abc	3661 abc	10139 bc	
Nonpareil-J	12803 abc		1.04 bcd	71.6 bcd	63.4 bcde	29.0 bc	3513 bc	9955 cd	
Winters	9434 ef		0.96 bcde	61.6 g	63.8 bcde	20.0 e	2415 e	9258 de	
Kahl	11035 cde		0.87 ef	59.1 g	79.2 a	21.1 de	2559 de	8513 ef	
Chips	9771 ef		0.93 def	58.6 g	55.9 de	20.0 e	2422 e	8144 ef	
Sweetheart	12798 abc		0.85 ef	73.3 abc	59.6 cde	24.0 d	2906 d	7965 fg	
Marcona	8977 fg		1.07 abc	32.5 h	77.7 a	21.2 de	2562 de	7563 gl	
Kochi	7252 g		1.17 a	68.9 de	52.6 e	18.7 e	2259 e	6955	
2010		Variety	No. of nuts/tree	Average kernel wt (g)	Shelling percentage	Kernel pounds per			Cumulative kernel yield (lbs/acre)
unit PAR int.	Tree	Acre							
Nonpareil-Nico	9521 abc		1.24 abcdef	72.5 ab	49.7 a	25.9 a	3141 a	14558 a	
Nonpareil-Newell	8429 cde		1.31 ab	73.6 a	45.2 abc	24.2 a	2931 a	14022 ab	
Nonpareil-3-8-2-70	8823 bcd		1.28 abcd	72.3 ab	47.1 ab	24.9 a	3011 a	13916 ab	
Nonpareil-Driver	8368 cde		1.28 abcd	71.0 ab	46.2 abc	23.6 a	2849 a	13911 ab	
Nonpareil-7	10612 ab		1.16 bcdef	69.8 ab	49.4 a	27.1 a	3282 a	13916 ab	
Nonpareil-5	9410 abc		1.24 abcde	72.3 ab	50.8 a	25.8 a	3130 a	13708 abc	
Nonpareil-6	9499 abc		1.21 abcdef	71.8 ab	48.7 ab	25.5 a	3081 a	13220 bc	
2-19e	6833 efg		1.10 bcdef	56.1 e	33.7 def	16.7 bc	2020 bc	13100 bc	
Nonpareil-Jones	8315 cde		1.23 abcdef	70.9 ab	43.8 abc	22.6 a	2737 a	12691 c	
Winters	6601 efg		1.11 bcdef	60.7 cde	38.5 bcde	16.0 bc	1945 bc	11203 d	
Chips	9089 abc		1.15 bcdef	65.9 abc	48.4 a	23.0 a	2789 a	10933 d	
Sweetheart	10915 a		0.80	71.8 ab	42.2 abcd	23.4 a	2839 a	10804 d	
Kahl	7587 cde		1.01	56.5 de	43.4 abcd	16.9 b	2048 c	10561 d	
Marcona	5073 gh		1.28 abc	26.2 g	36.7 cdef	14.4 bc	1745 bc	9308 e	
Kochi	3902 h		1.40 a	64.4 bcd	23.5 g	12.1 bc	1466 bc	8421 e	
2011		Variety	No. of nuts/tree	Average kernel wt (g)	Shelling percentage	Kernel pounds per			Cumulative kernel yield (lbs/acre)
unit PAR int.	Tree	Acre							
Nonpareil-Nico	18776 a		0.99 bcde	68.0 abc	90.1 a	41.0 a	4964 a	19523 a	
Nonpareil-3-8-2-70	17744 abc		1.05 bc	70.7 a	79.5 ab	41.0 a	4962 a	18878 ab	
Nonpareil-Newell	17790 abc		1.00 bcd	70.1 ab	69.4 abc	39.2 a	4745 a	18767 abc	
Nonpareil-Driver	17943 ab		0.98 bcde	66.0 abcd	72.8 ab	38.7 a	4683 a	18593 abc	
Nonpareil-7	17078 abcd		0.83 e	69.2 abc	72.3 ab	31.4 a	4555 a	18443 abc	
Nonpareil-5	15744 de		1.03 bc	70.4 ab	70.2 abc	35.9 a	4342 a	18050 bcd	
Nonpareil-6	16630 bcde		1.04 bc	70.0 ab	74.7 ab	38.2 a	4619 a	17838 bcd	
2-19e	18253 ab		0.91 bcde	64.8 abcd	65.2 bcd	36.8 a	4460 a	17560 cd	
Nonpareil-Jones	16993 abcd		0.96 bcde	70.0 ab	70.0 abc	36.0 a	4360 a	17051 d	
Winters	15979 cde		0.83 e	58.7 ef	56.8 cde	29.4 b	3554 b	14757 e	
Sweetheart	14969 e		0.86 de	64.1 bcde	47.3 cde	28.2 bc	3412 bc	14215 e	
Chips	11901 f		0.94 bcde	60.3 de	44.7 de	24.7 bcd	2985 bcd	13918 e	
Kahl	12420 f		0.89 cde	53.5 f	45.6 de	24.4 bcd	2953 bcd	13514 e	
Marcona	9633 g		1.07 b	30.8 c	45.0 de	22.7 d	2746 d	12054 f	
Kochi	8701 g		1.22 a	63.5 cde	9.4 e	23.3 d	2825 d	11247 f	
2012		Variety	No. of nuts/tree	Average kernel wt (g)	Shelling percentage	Kernel pounds per			kernel yield (lbs/acre)
unit PAR int.	Tree	Acre							
Nonpareil-Nico	9520 b		1.13 de	67.7 bcd	75.0 ab	23.6 a	2861 a	22384 a	
Nonpareil-3-8-2-70	8530 b		1.20 bc	70.9 bc	75.6 ab	22.6 ab	2733 ab	21611 ab	
Nonpareil-Newell	8481 b		1.15 cde	66.9 bcd	76.8 a	21.2 abc	2563 abc	21329 ab	
Nonpareil-Driver	8606 b		1.18 bcd	67.6 bcd	73.8 ab	22.3 ab	2695 ab	21288 ab	
Nonpareil-7	9262 b		1.14 cde	85.2 a	76.7 a	23.2 a	2811 a	21254 abc	
Nonpareil-5	8090 bc		1.19 bcd	69.0 bcd	74.1 ab	21.2 abc	2563 abc	20613 bc	
Nonpareil-6	11507 a		0.94	59.6 cd	69.3 c	23.8 a	2881 a	20441 bc	
2-19e	7617 bc		1.19 bcd	69.4 bcd	75.7 ab	20.1 abc	2432 abc	20270 bc	
Nonpareil-Jones	8855 b		1.18 bcd	67.7 bcd	73.3 b	23.0 ab	2783 ab	19833 c	
Winters	8679 b		1.01	61.9 bcd	61.0 d	19.3 abc	2338 abc	17095 d	
Sweetheart	8653 b		1.10 ef	59.8 cd	68.5 c	21.0 abc	2538 abc	16456 d	
Chips	9008 b		0.92	75.3 ab	76.2 ab	18.2 bc	2201 bc	16416 d	
Kahl	8830 b		1.05 fg	55.0 d	57.5 e	20.4 abc	2465 abc	15979 d	
Marcona	6449 c		1.22 b	65.5 bcd	74.5 ab	17.4 c	2104 c	13351 e	
Kochi	2025 d		1.41 a	26.0 e	60.0 de	6.3 d	763 d	12816 e	


**Table 4.** Number of nuts per tree, average kernel weight, shelling percentage, kernel pounds per unit of photosynthetically active radiation (PAR) intercepted, kernel pound per acre, cumulative kernel pounds per acre for the 2013 and 2014 seasons. Data for each year is sorted by cumulative yield. Selection 2-19e was recently released as Kester.

2013 Variety	No. of nuts/tree	Average kernel wt (g)	Shelling percentage	Kernel pounds per			Cumulative kernel yield (lbs/acre)
				unit PAR int.	Tree	Acre	
Nonpareil-Nico	20367 a	0.87 b	63.5 a	61.6 a	39.2 a	4738 a	27121 a
Nonpareil-3-8-2-70	18718 b	0.87 b	63.5 ab	56.5 ab	36.0 a	4354 a	25965 ab
Nonpareil-Newell	19539 ab	0.87 b	63.5 ab	58.0 ab	37.6 a	4545 a	25874 ab
Nonpareil-Driver	19539 ab	0.87 b	63.5 ab	60.2 a	37.4 a	4529 a	25817 ab
Nonpareil-7	19439 ab	0.87 b	63.5 ab	58.6 ab	37.4 a	4522 a	25776 ab
Nonpareil-5	18202 b	0.87 b	63.5 ab	55.4 ab	35.0 a	4234 a	24847 bc
Nonpareil-6	18769 b	0.87 b	63.5 ab	56.8 ab	36.1 a	4366 a	24636 bc
Nonpareil-Jones	18241 b	0.87 b	63.5 ab	54.7 ab	35.1 a	4243 a	24076 c
2-19e	16267 c	0.66 c	56.6 b	44.0 cd	23.9 c	2890 c	22958 c
Winters	13894 d	0.86 b	55.6 b	50.3 bc	26.5 bc	3201 bc	20296 d
Kahl	15587 c	0.85 b	55.3 b	57.9 ab	29.1 b	3524 b	19503 d
Chips	12689 d	0.89 b	57.3 b	39.7 de	24.9 bc	3010 bc	19466 d
Sweetheart	13943 d	0.78 b	66.5 a	40.4 de	24.0 c	2902 c	19318 d
Marcona	10858 e	1.11 a	28.9 d	49.4 bc	26.5 bc	3206 bc	16023 e
Kochi	7911	1.09 a	63.7 ab	33.5 d	19.0 d	2300 d	15651 e


2014 Variety	No. of nuts/tree	Average kernel wt (g)	Shelling percentage	Kernel pounds per			Cumulative kernel yield (lbs/acre)
				unit PAR int.	Tree	Acre	
Nonpareil-Nico	15387 b	0.91 de	70.1 bc	57.0 b	37.0 ab	4476 ab	31379 a
Nonpareil-3-8-2-70	15105 b	1.12 bcd	70.5 bc	57.6 b	37.4 ab	4522 ab	30486 ab
Nonpareil-Newell	14172 bc	1.12 bcd	69.8 bc	51.9 bc	34.8 ab	4208 ab	30082 abc
Nonpareil-Driver	13082 bcd	1.15 bc	70.3 bc	51.2 bc	33.1 b	4007 b	29824 bcd
Nonpareil-7	14201 bc	1.12 bcd	70.4 bc	52.1 bc	34.9 ab	4224 ab	29467 bcd
Nonpareil-5	14523 bc	1.15 bc	71.7 b	57.6 b	36.8 ab	4450 ab	29297 bcd
Nonpareil-6	13930 bc	1.13 bcd	69.5 bcd	52.8 bc	34.7 ab	4195 ab	28831 cd
Nonpareil-Jones	16267 c	0.86 e	57.6 d	48.4 bc	32.2 b	3901 b	28584 d
2-19e	14378 bc	0.92 cde	69.5 bcd	49.1 bc	29.1 bc	3616 bc	27075 e
Winters	17427 a	0.91 de	77.1 a	67.0 a	35.0 ab	4237 ab	24532 f
Chips	11188 cde	0.98 cde	67.7 bcd	39.9 cd	23.9 cd	2886 cd	22353 g
Kahl	9310 e	1.03 bcde	55.3 e	40.6 cd	21.0 d	2543 d	22046 g
Sweetheart	10145 de	0.92 cde	66.7 cd	31.5 d	20.5 d	2477 d	21794 g
Kochi	5981 f	1.25 b	66.8 cd	26.4 d	16.5 d	1996 d	17646 h
Marcona	1721 g	1.48 a	57.6 e	10.3 e	5.6 e	682 e	16705 h

**Table 5.** Varieties and selections planted at the next generation regional almond variety trials. Items 1-31 are planted at all 3 sites while additional material planted at individual sites is listed at the end. Trees at the Butte, Stanislaus and Madera sites were planted on Krymsk 86, Nemaguard and Hansen 536 rootstock respectively (exceptions are noted at bottom of table).


	Variety	Source	Provides Bud wood
1	Eddie	Bright's	Bright's
2	Capitola	Burchell	Burchell
3	Supareil	Burchell	Burchell
4	self-fruitful P16.013	Burchell	Burchell
5	Self-fruitful P13.019	Burchell	Burchell
6	Booth	Burchell	Burchell
7	Sterling	Burchell	Burchell
8	Bennett	Duarte	Duarte
9	Nonpareil	Fowler	Fowler
10	Durango	Fowler	Fowler
11	Jenette	Fowler	Fowler
12	Aldrich	Fowler	Fowler
13	Marcona	Spain	Gradziel
14	Winters	UCD	Fowler
15	Sweetheart	UCD	Fowler
16	2-19e (Kester)	UCD	Gradziel
18	UCD3-40	UCD	Gradziel
19	UCD18-20	UCD	Gradziel
20	UCD1-16	UCD	Gradziel
21	UCD8-160	UCD	Gradziel
22	UCD8-27	UCD	Gradziel
23	UCD1-271	UCD	Gradziel
24	UCD1-232	UCD	Gradziel
25	UCD7-159	UCD	Gradziel
26	UCD8-201	UCD	Gradziel
27	Y121-42-99	USDA	Ledbetter
28	Y117-86-03	USDA	Ledbetter
29	Y116-161-99	USDA	Ledbetter
30	Y117-91-03	USDA	Ledbetter
31	Folsom	Wilson	Wilson
Chico site only			
Wood Colony on Krymsk 86			
Chowchilla site only			
32	Lone Star on Hansen	Burchell	Burchell
Stanislaus site only			
2-19e (Kester) on Nemaguard and Hanson			




**Figure 1.** Average yield for all varieties, selections and Nonpareil sources by orchard age for the 1993 Butte, Delta and Kern Regional Variety Trials as well as the McFarland Variety Trial that was planted in 2004. Number in parenthesis after trial name is number of trees per acre.




**Figure 2.** Bloom data for the McFarland Trial for the 2010-2012 seasons. Cross hatched area indicates onset of bloom to 100% petal fall. “F” indicates full bloom.




**Figure 3.** Bloom data for the McFarland Trial for the 2013 and 2014 seasons. Cross hatched area indicates onset of bloom to 100% petal fall. "F" indicates full bloom.




**Figure 4.** Average bloom data (2006 to 2014) for the McFarland Trial. Cross hatched area indicates onset of bloom to 100% petal fall. "F" indicates full bloom.




**Figure 5.** Hull split progression for the McFarland Trial for the 2010-2013 seasons. Cross hatched area indicates onset of bloom to 100% petal fall. "F" indicates full bloom.



**Figure 6.** Hullsplit progression by variety and Nonpareil source for 2014 season at the McFarland trial.



**Figure 7.** Average hullsplit progression by variety and Nonpareil source for 2006 to 2014 seasons at the McFarland trial.



**Figure 8.** Yield per unit PAR intercepted by variety and year (kernel pounds per 1% light intercepted). Number following variety name in legend indicates the five year average value. Average for all varieties and years was 49.3. A value of 50 is what we have found the best orchards can average over several years.