

MATING DISRUPTION OF THE NAVEL ORANGEWORM

Ring T Cardé, Brad Higbee¹ and Robbie Girling²

Department of Entomology, University of California, Riverside CA, 92521 USA

¹ Paramount Farms. ² University of Southampton (U.K.)



Objectives:

1. Compare in a wind-tunnel assay the in-flight and close-range male orientation and courtship behaviors evoked by point-source formulations of pheromone, including dispensers prepared for evaluation in the field. Determine which blend of components evokes the highest proportion of source finding.
2. Use the wind tunnel-assay to determine why some batches of synthetic pheromone and formulated lures are more attractive than others. In collaboration with Brad Higbee, field test formulated dispensers of pheromone for efficacy as lures in traps.
3. Determine the role of sound in courtship.
4. Determine the pattern of pheromone dispersal, with particular attention to vertical movement of odor plumes, using the almond orchard as a model habitat. These measurements will aid in determining the optimal height in the canopy for placement of monitoring traps and puffers.

Interpretive Summary:

We have two overarching goals: 1) to improve the efficacy of mating disruption in the navel orangeworm moth, *Amyelois transitella*, referred to as “NOW.” 2) To aid in the development of a highly attractive lure that would be useful as a monitoring tool in pest management programs. To these ends we have: 1) Optimized the composition and ratio of components for maximal male attraction in a wind tunnel using a simple, filter-paper dispenser. 2) Shown that four expected breakdown products of the main aldehyde pheromone component do not affect attraction. 3) Tested a number of lures formulated to provide longevity in the field and competitiveness with female-baits. 4) Documented the courtship behaviors and sensory inputs (pheromone, sound and tactile) that occur once the male and female are in proximity. 5) Established the patterns of pheromone dispersal in almond orchards using visual tracers and measured wind flow and turbulence patterns with a sonic 3-D anemometer.

There are a number of potential components of the pheromone (Leal et al. 2005). One of our goals has been to verify which of the 9 published compounds (as well as others newly discovered but unpublished) mediate attraction and courtship. We have now defined an optimal blend of 4 components (Kanno et al. 2010; Kuenen et al. 2010). Such information is a crucial step in devising highly attractive lures for monitoring traps. Furthermore, it is widely thought that the complete pheromone blend should be the most efficacious mixture for mating disruption (Minks and Cardé 1988, Cardé and Minks 1995; Cardé 2007).

Results and Discussion:

1. Upwind flight along the pheromone plume and landing on the odor source required the simultaneous presence of two components, (11Z,13Z)-hexadecadienal, (3Z,6Z,9Z,12Z,15Z)-tricosapentaene, and the addition of either (11Z,13Z)-hexadecadien-1-ol or (11Z,13E)-hexadecadien-1-ol. A mixture of all four components produced the highest levels of rapid source location and source contact. In wind-tunnel assays, males did not seem to distinguish among a wide range of ratios of any of the three components added to (11Z,13Z)-hexadecadienal. Dosages of 10 and 100 ng of the 4-component blend produced higher levels of source location than dosages of 1 and 1000 ng (Kanno et al. 2010; Kuenen et al. 2010). The broad range of component ratios that evoke attraction will simplify some aspects of development of a field lure.

2. We are working with Walter Leal and Jocelyn Millar is trying to identify contaminants and breakdown products that may compromise the efficacy of lures in the field. So far four possible contaminants have been evaluated for their possible suppressive effect on attraction. These compounds were prepared and supplied courtesy of Bedoukian Research. Briefly, adding any of the three geometrical isomers of the aldehyde pheromone component (the 11Z, 13E, the 11E, 13Z, and the 11E, 13E) pheromone component to the complete blend did not diminish attractiveness. Work with many other moth species often has found that geometrical isomers of monounsaturated or doubly unsaturated pheromones are inhibitory to attraction, but this is not the case with the NOW. Similarly, the adding the acid of the aldehyde to the 4-component blend also did not affect attractiveness. This is very helpful to lure formulation, in that such breakdown products would be expected.

A second approach has been to compare, for example, lures using the principal pheromone component (the aldehyde) from different sources, or following purification, while holding the other 3 components constant. Clearly batches have differed somewhat in their attractiveness, but the chemical identity(ies) of the antagonist(s) remains unclear.

Work on lures for the field has relied on membrane dispensers prepared by Suterra. These have been evaluated in the field by Brad Higbee and in our wind tunnel. Our tests used dispensers aged outside in traps between wind-tunnel evaluations. A few dispenser and synthetic combinations have proved very encouraging. For example, two of these were evaluated with 360 males each over 50 days. One formulation evoked 31.1% lure contact and another evoked 35.8% lure contact; both formulations appeared to be as attractive at the end of the test as at the beginning.

3. Courting males produce ultrasonic “clicks” during wing fanning. This signal inhibits calling (pheromone-emitting) females from moving away when contacted by a male and increases the likelihood of mating. This study is complete and is being prepared for publication.

4. Nighttime observations in almond orchards by Brad Higbee indicated that pheromone plumes might have considerable vertical movement. Together, we documented this dispersal pattern, initially using titanium tetrachloride sources positioned from a meter above ground level to the top of the canopy.

This past field season we documented wind flow in 3-D with a sonic anemometer at 6 heights from 2.08 to 6.65 meters above the ground level, monitoring the speed, turbulence and direction of airflow for 5 minutes at each height every 3 hours over 4 days. We are now analyzing these recordings for publication, but we provide some comparative data (Fig. 1). These records show that the net flow of air during the time when NOW mate (late night to dawn) is upward and not simply planar as has been generally assumed.

One practical implication of these observations for mating disruption is that when pheromone puffers are deployed at height of the top of the orchard canopy, as is current practice, much of their output may be carried upwards beyond the canopy. This suggests that much of the output of puffers so deployed would be unavailable to disrupt NOW mate finding—that it would be “wasted.” We suggest that puffers would be just as efficacious if deployed at mid or lower canopy levels. As a result of these observations and his own work, Brad Higbee has evaluated male capture in female-baited traps positioned at several canopy heights and the effect of the height of puffers on the efficacy of mating disruption.




Fig.1. Percentages of time sorted into 10° bins of wind deviating from horizontal (= 0°) during 5-minute-long samples recorded at 00:00 (midnight), 03:00, 06:00 and 09:00 at the indicated heights. The 6.65 m height is more than a meter above the canopy’s highest point. Although there is considerable flow in the horizontal plane, at nighttime when NOW is sexually active, clearly much of the flow is upward. Wind velocities (not shown) are attenuated within the canopy.

Research Effort Recent Publications:

Girling, R.D. and R.T. Cardé. 2006. Analysis of the courtship behavior of the navel orangeworm, *Amyelois transitella* (Walker) (Lepidoptera: Pyralidae), with a commentary on methods for the analysis of sequences of behavioral transitions. *J. Insect Behav.* 19:497-520.
Kanno, H., L.P.S. Kuenen, K.A. Klingler, J.G. Millar and R.T. Cardé. 2010. Attractiveness of a four-component pheromone blend to male navel orangeworm moths. *J. Chem. Ecol.* 36:584-591.
Wang, H.-L., C.-H. Zhao, J. G. Millar, R.T. Cardé and C. Löfstedt. 2010. Biosynthesis of unusual moth pheromone components involves two distinctly different pathways in the navel orangeworm, *Amyelois transitella*. *J. Chem. Ecol.* 36:535-547.

References Cited:

Cardé, R.T. 2007. Using pheromones to disrupt mating of moth pests. In: *Perspectives in Ecological Theory and Integrated Pest Management*, M. Kogan and P. Jepson (eds.) Cambridge University Press, Cambridge, pp. 122-169.
Cardé, R.T. and A.K. Minks. 1995. Control of moth pests by mating disruption: successes and constraints. *Annu. Rev. Entomol.* 40:559-585.
Kanno, H., L.P.S. Kuenen, K.A. Klingler, J.G. Millar and R.T. Cardé. 2010. Attractiveness of a four-component pheromone blend to male navel orangeworm moths. *J. Chem. Ecol.* 36:584-591.
Kuenen, L.P.S., J.S. McElfresh and J.G. Millar. 2010. Identification of critical secondary components of the sex pheromone of the navel orangeworm, (Lepidoptera: Pyralidae). *J. Econ. Entomol.* 103:314-330.
Leal, W.S., A.L. Parra-Pedrazzoli, K.-E. Kaissling, T.L. Morgan, F.G. Zalom, D.J. Pesak, E.A. Dundulis, C.S. Burks and B.S. Higbee. 2005. Unusual pheromone chemistry in the navel orangeworm: novel sex attractants and a behavioral antagonist. *Naturwissenschaften* 92: 139-146.
Minks, A.K. and R.T. Cardé. 1988. Disruption of pheromone communication in moths: is the natural blend really most efficacious? *Entomol. Exp. Appl.* 49:25-36.

Acknowledgments—Wind tunnel experiments were conducted mainly by Kim Klingler. Jocelyn Millar (UCR) provided most of the test chemicals. Brad Higbee (Paramount Farms) conducted field trapping tests. Work on the pattern of pheromone dispersal was a joint with Brad Higbee and Robbie Girling (Southampton University). Additional collaborators include Walter Leal (UC Davis), Bedoukian Research and Suterra. Equal support from the California Pistachio Commission is gratefully acknowledged.