

Effects of *Xylella fastidiosa* Group, Almond Cultivar, and Climate on the Establishment and Persistence of Almond Leaf Scorch

Project No.: 06-PATH5-Daane

Project Leader: Kent Daane
Environmental Science
Policy and Management Department
137 Mulford Hall
University of California
Berkeley CA 94720-3114
daane@ucl.ac.uk

Project Cooperators and Researchers:

Christina Wistrom: ESPM Dept., UC Berkeley.
Bruce Kirkpatrick, Plant Pathology, Dept., UC Davis.

Objectives:

1. Compare establishment and over wintering of grape and almond strain *X. fastidiosa* in susceptible and resistant almond cultivars, and
2. Determine the effects of cold treatment on the over winter survival of *X. fastidiosa* in almonds.

Interpretive Summary:

Controlled temperature study: Five months after inoculation with ALS 6 *X. fastidiosa*, 72% of inoculated trees (89 of 125) developed almond leaf scorch, averaging 12.8 (SE = 1.25) symptomatic leaves per infected tree, and a median population of 4.4×10^6 CFU/g. None of the buffer-inoculated trees were infected with *X. fastidiosa*, and they averaged 0.3 scorched or yellowed leaves per tree. Fourteen of 89 infected trees died during or after cold treatment, compared to 1 of 27 buffer-inoculated trees (χ^2 with Yates' correction = 3.17; $P > 0.05$; $df = 1$). Six of 29 trees were negative for *X. fastidiosa* after one month cold treatment, compared to 4 of 30 after two months, and 2 of 30 after four months (**figure 1A**). There were not significant differences in the number of recovered trees regardless of temperature or time (χ^2 with pairwise comparisons).

ALS symptoms were twice as severe after cold treatment as before. In 2005, buffer-inoculated trees averaged 0.26 scorched or yellowed leaves per tree (SE = 0.25), compared to 12.8 symptomatic leaves per *X. fastidiosa*-infected tree (SE = 1.23). In 2006, there were an average of 1.29 (SE = 0.42) scorched and yellowed leaves per buffer inoculated tree and 30.24 (SE=2.57) symptomatic leaves per *X. fastidiosa* infected tree. Symptoms were worse in trees following four months of cold treatment, compared to trees with one and two months (**figure 1B**).

The time of dormancy, but not temperature, influenced symptom severity (Time $P = 0.0002$, Temperature $P = 0.593$, Temperature*Time $P = 0.39$). *X. fastidiosa* in infected trees were similar across all treatments (2-way ANOVA, Time $P = 0.74$; Temperature $P = 0.19$; Time*Temperature Interaction $P = 0.39$) with a median population of 2.82×10^6 CFU/g. Trees in the 1.7°C and 7°C treatments were subjected to temperatures below the growth threshold of *X. fastidiosa* for the entire period: 769 hours for 1 month, 1441 hours for 2 months, and 2905 hours for 4 months. Outside, trees were below 7°C for 262 hours in the 1 month treatment, 557 hours in the 2 month treatment, and 673 hours for the 4 month treatments. Only trees kept outside were subjected to sub-freezing temperatures, for 10 hours, both in the 2 and 4 month treatments.

Field sites: In 2005, two to four months after inoculation, *X. fastidiosa* was recovered from 49% of bacteria-inoculated trees (76 of 155), and from none of the buffer-inoculated trees. There was no difference in the proportion of infected trees at UCD or at IRC (36 of 76 at UCD; 40 of 79 at IRC; $\chi^2 = 0.17$; $P > 0.05$; $df = 1$; figure 4), nor in median *X. fastidiosa* populations in infected trees (4.0×10^6 colony-forming units per gram of petiole tissue at UCD; 1.2×10^7 CFU/g at IRC; t-test with \log_{10} -transformed data; $P = 0.18$; $df = 75$). However, disease symptoms were more severe at UCD, especially in 'Peerless' trees, averaging 8.8 (SE = 2.1) symptomatic leaves per infected tree, compared to 2.3 leaves per infected 'Butte' tree (SE = 0.7; $P = 0.02$; two-sample t-test; $df = 35$). At IRC, symptoms were negligible, as both varieties averaged 0.3 scorched leaves per infected tree (SE = 0.1; $P = 0.76$; two-sample t-test; $df = 39$). Background yellowing and scorching in uninfected trees was 0.11 leaves per tree (SE = 0.08) at UCD and 0.21 (SE = 0.1) leaves per tree at IRC (**Figure 2A**).

Grape strain *X. fastidiosa* infected trees more frequently than almond *X. fastidiosa* at UCD but not at IRC (**figure 2B**). At UCD, 4 of 37 trees inoculated with Dixon or ALS-6 developed infections, compared to 32 of 39 trees with Fresno-ALS or Medeiros (χ^2 with Yates' correction = 38.71; $P < 0.001$; $df = 1$). At IRC, both almond and grape strains of *X. fastidiosa* infected trees with the same frequency (16 of 39 inoculated with almond; 24 of 40 inoculated with grape strain; $\chi^2 = 2.84$; $P > 0.05$; $df = 1$). Grape and almond strains reached similar titers in infected plants, median 3.48×10^6 CFU/g for almond strain *X. fastidiosa*, and 5.71×10^6 CFU/g for grape strain, both sites combined. All *X. fastidiosa* recovered from inoculated trees matched the type initially inoculated; there was no *X. fastidiosa* movement between trees at either field site.

Over the winter, two trees died at UCD, and 62 trees or inoculated branches died at IRC. Surviving trees (previously infected in 2005) at IRC were evenly distributed among PD, ALS and buffer isolate treatments, with 6 buffer-inoculated, 3 ALS 6-inoculated, 4 Dixon-inoculated, 4 Fresno-inoculated, and 6 Medeiros-inoculated trees surviving. While mortality was high, similar losses were seen in previous studies examining the over winter survival of *X. fastidiosa* in grapevines in extremely cold climates (Purcell 1980). No *X. fastidiosa* was recovered from trees at IRC in 2006. At UCD, *X. fastidiosa* was recovered from only one previously-infected tree, the Medeiros isolate in a 'Peerless' tree. At both sites, there was negligible leaf scorch and chlorosis in uninfected trees. 'Butte' trees at IRC were beginning to senesce at the time of assessment. Trees at UCD were subjected to 1076 hours of temperatures below 7°C between inoculation and

rating in 2006 (1070 over winter), including 44 hours below 0°C. (An average of 2928 hours elapsed between inoculation and rating in 2005, and 12,223 hours in 2006). Trees at IRC received four times as much cold, 4659 hours over winter between inoculation in July 2005 and rating for disease in September 2006 (4600 over winter) of 10,343 total hours. Trees spent 1852 hours below 0°C at IRC.

Implications for control. The prevalence of almond-strain *X. fastidiosa* in northern almond orchards may be explained if almond strains initially infect trees at low rates but survive the winter more frequently than grape strains. Data from this study supported half this hypothesis, since almond strains initially infected 11 to 15% of trees, similar to the 21 to 33% infection rate previously reported by other researchers. Grape *X. fastidiosa* isolates infected 79% of inoculated trees at UCD in this study, and 64 and 75% in field trials in Davis and Parlier. In this study, no almond infections and only one grape strain *X. fastidiosa* infections overwintered in field plots, although almond-strain *X. fastidiosa* overwintered in potted plants.

Since only one infection survived the winter in the field plots, there is so far no data to support the hypothesis that *X. fastidiosa* infections overwinter more frequently in susceptible 'Peerless' compared to 'Butte' since all but one *X. fastidiosa* infections died over winter. The one surviving infection was in susceptible 'Peerless'. Repeated inoculations at UCD in 2007 will provide more data to test this hypothesis. The effect of cold on *X. fastidiosa* survival was previously investigated in grapes but not in almonds. Exposure to very cold overwintering temperatures can eliminate *X. fastidiosa* infections but also results in significant plant mortality. Potted almond trees in the controlled study exposed to 4-month dormancy had more symptomatic leaves than trees dormant for one or two months. This is in contrast to previous studies in grapes where *X. fastidiosa* populations decreased 320-fold in only 18 days at 5°C (41°F). A second replication of the cold-chamber experiment is ongoing to test the hypothesis that threshold temperatures to kill almond-strain *X. fastidiosa* are lower than those needed to kill grape *X. fastidiosa* in almond trees.

Funding. This research was jointly funded by the Almond Board of California (2005-2006) and the University of California Pierce's Disease and Glassy-winged Sharpshooter Program (2006-2007). The proceedings article provided here is a summary for almond growers for the 2005-2006 Almond Board of California funded program.

Figure 1. A) *X. fastidiosa* infections and B) Almond Leaf Scorch symptoms in previously-infected potted almond trees after 1, 2 or 4 months of dormancy at 1.7°C, 7°C, or outside in Parlier, CA.




Figure 2. A) Almond leaf scorch symptoms and B) *X. fastidiosa*-infections in 'Butte' and 'Peerless' almond trees at Davis (left) and Tulelake (right) field sites 2-4 months after inoculation.

