

Quantifying Varroa Resistance to Miticides in US Honey Bee Colonies

Project Leader: Dennis vanEngelsdorp

CMNS – Entomology, University of Maryland, College Park, MD 20742
(717) 884-2414; dennis.vanengelsdorp@gmail.com

PROJECT SUMMARY

Objectives:

- Survey the susceptibility of the US varroa mite population to both registered and unregistered mite control products.
- Rapidly communicate mite susceptibility and resistance levels to US beekeepers so they can make informed mite control decisions.

Background and Discussion:

While many factors contribute to honey bee declines, there is consensus that the single most important contributing factor is varroa mite. To control these parasites, beekeepers rely on a handful of mite control products. Unfortunately, mites have developed resistance to many of these products, and/or the product’s efficacy is questionable. This is of serious concern, as beekeepers need to be aware of which products will be effective. With some assurance as to which products will work, beekeepers can avoid unnecessary treatments, reduce unexpected losses, and improve the general welfare of their colonies. Further, there is a critical need for resistance management. One aspect of this is to bring to the market novel varroacide chemistries that target unique resistance avoidance pathways.

We began performing resistance assays on live bees collected for National Honey Bee disease survey in April 7, 2014 and for the BIP tech teams collected samples beginning August 2014. In all samples from 365 operations were received with sufficient bees to perform a resistance assay using one or more mite control product. The resistance assay involves placing approximately 300 bees in a cage along with a small strip of miticides (after Pettis et al. 1998). Mites that fell from the bees after 6 hours exposure were

counted. Sugar powder was used to remove and quantify any mites remaining on the caged bees. When the total number of mites removed from caged bees was less than 5, the assay was not considered for further analysis. Samples from 219 different operations were collected that has sufficient mites to perform at least one successful assay.

A total of 4 different products were tested for resistance (Table 1), all but flumethrin are labeled for use by beekeepers in the US. A population of mites was considered resistant, if the miticide failed to kill less than 20% of the mites and was considered susceptible if it killed 80% or more of the mites in a sample.

There was no evidence of amitraz resistance in tested colonies, but only 20% of populations were considered truly susceptible (**Table 1**). Comaphos resistance was evidenced in 12% of tested populations, while 21% of the mite population was considered susceptible. Fluvalinate resistance was wide spread with 72% of tested populations being resistant and no population being susceptible. 3% of flumethrin tested mite populations were resistant while 47% of populations were susceptible. These results reaffirm the urgent need for additional mite control products for the bee industry.

Table 1: Percentage of mite populations with sufficient mites to permit testing that were resistant or highly resistant to mites.

Product	n	Resistant (%)	Susceptible (%)
Amitraz	176	0	20
Coumaphos	169	12	21
Flumethrin	114	3	47
Fluvalinate	174	72	0

Project Cooperators and Personnel: Sam Abban and Jeff S. Pettis, USDA-ARS, Beltsville Bee Lab; Troy Anderson, Virginia Tech.

For More Details, Visit

- Poster location 10, Exhibit Hall A + B during the Almond Conference; or on the web (after January 2016) at Almonds.com/ResearchDatabase
- 2014 - 2015 Annual Reports CD (14-POLL6-vanEngelsdorp); or on the web (after January 2016) at Almonds.com/ResearchDatabase
- Related project: 15-POLL17-Anderson